

BERGMAN KERNELS AND EQUILIBRIUM MEASURES FOR LINE BUNDLES OVER PROJECTIVE MANIFOLDS

ROBERT BERMAN

ABSTRACT. Let L be a holomorphic line bundle over a compact complex projective Hermitian manifold X . Any fixed smooth hermitian metric ϕ on L induces a Hilbert space structure on the space of global holomorphic sections with values in the k th tensor power of L . In this paper various convergence results are obtained for the corresponding Bergman kernels (i.e. orthogonal projection kernels). The convergence is studied in the large k limit and is expressed in terms of the equilibrium metric ϕ_e associated to the fixed metric ϕ , as well as in terms of the Monge-Ampere measure of the metric ϕ itself on a certain support set. It is also shown that the equilibrium metric is $\mathcal{C}^{1,1}$ on the complement of the augmented base locus of L . For L ample these results give generalizations of well-known results concerning the case when the curvature of ϕ is globally positive (then $\phi_e = \phi$). In general, the results can be seen as local metrized versions of Fujita's approximation theorem for the volume of L .

CONTENTS

1. Introduction	1
2. Preliminaries: positivity and base loci	8
3. Equilibrium measures for line bundles	11
4. Bergman kernel asymptotics	22
5. Examples	32
References	34

1. INTRODUCTION

Let L be a holomorphic line bundle over a compact complex projective manifold X of dimension n . Fix a smooth Hermitian fiber metric, denoted by ϕ , on L and a smooth volume form ω_n on X . The curvature form of the metric ϕ may be written as $dd^c\phi$ (see section 1.4 for definitions and further notation). Denote by $\mathcal{H}(X, L^k)$ the Hilbert space obtained by equipping the space $H^0(X, L^k)$ of global holomorphic sections with values in the tensor power L^k with the norm induced by the given metric ϕ on L and the volume form ω_n . The *Bergman kernel* of the Hilbert space $\mathcal{H}(X, L^k)$ is the integral kernel of the orthogonal projection from the space of all smooth sections with values in L^k onto $\mathcal{H}(X, L^k)$. It may be represented

Key words and phrases. Line bundles, holomorphic sections, Bergman kernel asymptotics, global pluripotential theory. *MSC (2000):* 32A25, 32L10, 32L20, 32U15.

by a holomorphic section $K_k(x, y)$ of the pulled back line bundle $L^k \boxtimes \overline{L}^k$ over $X \times \overline{X}$ (formula 4.1).

In the case when the curvature form $dd^c\phi$ is globally *positive* the asymptotic properties of the Bergman kernel $K_k(x, y)$ as k tends to infinity have been studied thoroughly with numerous applications in complex geometry and mathematical physics. For example, $K_k(x, y)$ admits a complete local asymptotic expansion in powers of k ; the Tian-Zelditch-Catlin expansion (see [37, 8] and references therein). The point is that when the curvature form $dd^c\phi$ is globally positive, the Bergman kernel asymptotics at a fixed point may be localized and hence only depend (up to negligible terms) on the covariant derivatives of $dd^c\phi$ at the fixed point.

The aim of the present paper is to study the case of a general smooth metric ϕ on an arbitrary line bundle L over a projective manifold, where global effects become important and where there appears to be very few previous general results even in the case when the line bundle L is ample (and even when X is a complex curve). We will be mainly concerned with three natural *positive* measures on X associated to the setup introduced above. In order to introduce these measures first assume that the line bundle L is ample. The first measure on X to be considered is the *equilibrium measure*

$$\mu_\phi := (dd^c\phi_e)^n/n!,$$

where ϕ_e is the *equilibrium metric* defined by the upper envelope 3.1 (i.e. $\phi_e(x) = \sup \tilde{\phi}(x)$, where the supremum is taken over all metrics $\tilde{\phi} \leq \phi$ with positive curvature). For example, when X is the projective line $\mathbb{P}^1$ and L is the hyperplane line bundle $\mathcal{O}(1)$ the measure μ_ϕ is a minimizer of the “weighted logarithmic energy” [32]. Next, the weak large k limit of the measures

$$(1.1) \quad k^{-n} B_k \omega_n,$$

where $B_k(x) := K_k(x, x)e^{-k\phi}$ will be referred to as the *Bergman function* is considered and finally the limit of the measure

$$(dd^c(k^{-1}\ln K_k(x, x)))^n/n!,$$

often referred to as the *kth Bergman volume form* on X associated to (L, ϕ) (and $k^{-1}\ln K_k(x, x)$ is called the *kth Bergman metric* on L).

When L is ample it is well-known that the integrals over X of all three measures coincide. In fact, the integrals all equal the integral over X of the possibly non-positive form $(dd^c\phi)^n/n!$, as is usually shown by combining the Riemann-Roch theorem with Kodaira vanishing. The main point of the present paper is to show the corresponding *local* statement. In fact, all three measures will be shown to coincide with the measure

$$1_D(dd^c\phi)^n/n!$$

where 1_D is the characteristic function of the set

$$(1.2) \quad D = \{\phi_e = \phi\} \subset X$$




FIGURE 1.1. In the toric case (see example 5.2) the metric ϕ is represented by a function on $\mathbb{R}^n$ and the graph of the equilibrium potential ϕ_e (the dashed line) is obtained as the *convex hull* of the graph of ϕ . The set D is then the projection of the set where the two graphs coincide.

In the case when the metric ϕ has a semi-positive curvature form, $\phi_e = \phi$, i.e. the set D equals all of X .

Before turning to the the statement of the main general results, note that when L is not ample, the main new feature is that the equilibrium metric ϕ_e will usually have *singularities* (i.e. points where it is equal to $-\infty$) and its curvature $dd^c\phi_e$ is a positive *current*. However, as is well-known it does give a metric on L with *minimal* singularities. Such metrics play a key role in complex geometry (compare remark 3.5). Similarly, the Bergman metric $k^{-1}\ln K_k(x, x)$ is singular along the base locus $\text{Bs}(|kL|)$ of L , i.e. along the common zero-locus of the sections in $H^0(X, L^k)$. Still, the convergence results referred to above in the case when L is ample will be shown to hold provided that the measures are extended by zero over the singularities. Then integrating over X gives new proofs of Boucksom's version of Fujita's approximation theorem for the *volume* of the line bundle L [22, 10] and its interpretation in terms of intersection of zero-sets of sections in $H^0(X, L^k)$ by Demailly-Ein-Lazarsfeld [20]. Finally, the asymptotic properties of the full Bergman kernel $K_k(x, y)$ are studied.

The present approach to the Bergman kernel asymptotics is based on the use of "local holomorphic Morse-inequalities", which are local version of the global ones introduced by Demailly [15]. These inequalities are then combined with some L^2 -estimates and global pluripotential theory, the pluripotential part being based on the recent work [24] by Guedj-Zeriahi. Conversely, it turns out that several basic, but non-trivial, results in pluripotential theory may obtained as consequences of the Bergman kernel asymptotics (compare for example remark 3.2).

A crucial step is to first show the $\mathcal{C}^{1,1}$ -regularity of the equilibrium metric ϕ_e on the complement of the augmented base-locus $\mathbb{B}_+(L)$, which should be of independent interest.

1.1. Statement of the main results. Assume that (L, ϕ) and (F, ϕ_F) are smooth Hermitian line bundles over X and denote by $E(k)$ the twisted line bundle $L^k \otimes F$. The equilibrium measure on X (associated to the smooth metric ϕ on L) is defined as the positive measure

$$\mu_\phi := 1_{U(L)}(dd^c\phi_e)^n/n!,$$

where $U(L)$ is the open set in X where ϕ is locally bounded (see section 3). The first theorem to be proved is used to express μ_ϕ in terms of $(dd^c\phi)^n$ on the set D (formula 1.2 above).

Theorem 1.1. *Suppose that L is a big line bundle and that the given metric ϕ on L is smooth (i.e. in the class $\mathcal{C}^2$). Then the equilibrium metric ϕ_e is locally in the class $\mathcal{C}^{1,1}$ on $X - \mathbb{B}_+(L)$ i.e. ϕ_e is differentiable and all of its first partial derivatives are locally Lipschitz continuous there. Moreover, the equilibrium measure satisfies*

$$\mu_\phi n! = 1_{X - \mathbb{B}_+(L)}(dd^c\phi_e)^n = 1_D(dd^c\phi)^n = 1_{D \cap X(0)}(dd^c\phi)^n$$

in the sense of measures, where $X(0)$ is the set where $dd^c\phi > 0$.

The regularity theorem is essentially optimal (compare the examples 5.2 and 5.3). The next theorem which should be considered as the main result of this paper gives that, in general, the measure $k^{-n}B_k\omega_n$ introduced above (formula 1.1) converges to the equilibrium measure μ_ϕ .

Theorem 1.2. *Let B_k be the Bergman function of the Hilbert space $\mathcal{H}(X, E(k))$. Then*

$$(1.3) \quad k^{-n}B_k(x) \rightarrow 1_{D \cap X(0)} \det(dd^c\phi)(x)$$

for almost any x in X , where $X(0)$ is the set where $dd^c\phi > 0$ and D is the set 3.3. Moreover, the following weak convergence of measures holds:

$$k^{-n}B_k\omega_n \rightarrow \mu_\phi,$$

where μ_ϕ is the equilibrium measure.

The Bergman function B_k may be interpreted as a “dimensional density” of the Hilbert space $\mathcal{H}(X, E(k))$. The asymptotic (normalized) dimension of $\mathcal{H}(X, L^k)$ is called the *volume* of a line bundle L [30]:

$$(1.4) \quad \text{Vol}(L) := \limsup_k k^{-n} \dim H^0(X, L^k)$$

Integrating the convergence of the Bergman kernel in the previous theorem now gives the following version of Fujita’s approximation theorem [22, 10] (compare remark 4.9 for a comparison with closely related expressions of $\text{Vol}(L)$).

Corollary 1.3. *The volume of a line bundle L is given by the total mass of the equilibrium measure:*

$$(1.5) \quad \text{Vol}(L) = \int_X \mu_\phi$$

and $\text{Vol}(L) = 0$ precisely when L is not big.

The following theorem gives, in particular, the weak convergence on X of the k th Bergman volume forms (extended by zero over the base-locus of $E(k)$).

Theorem 1.4. *Let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, E(k))$. Then the following convergence of Bergman metrics holds:*

$$k^{-1}\phi_k \rightarrow \phi_e$$

uniformly on any fixed compact subset Ω of $X - \mathbb{B}_+(L)$. More precisely,

$$(1.6) \quad e^{-k(\phi - \phi_e)} C_\Omega^{-1} \leq B_k \leq C_\Omega k^n e^{-k(\phi - \phi_e)}$$

Moreover, the corresponding k th Bergman volume forms converge to the equilibrium measure:

$$1_{X - \text{Bs}(|E(k)|)} (dd^c(k^{-1} \ln K_k(x, x)))^n / n! \rightarrow \mu_\phi$$

weakly as measures on X .

The weak convergence in the theorem above on $X - \mathbb{B}_+(L)$ is a consequence of the uniform convergence of the Bergman metrics $k^{-1}\phi_k$ on compacts of $X - \mathbb{B}_+(L)$. But to get the weak convergence on all of X theorem 1.2 (or rather its corollary 1.3) is invoked.

For an ample line bundle L it is a classical fact that the volume $\text{Vol}(L)$ may be expressed as an intersection number L^n . More generally, for any line bundle L over X the intersection of the zero-sets of n “generic” sections in $H^0(X, L^k)$ with $X - \text{Bs}(|kL|)$ (the complement of the common zero-locus of all sections) is a finite number of points. The number of points is called the *moving intersection number* and is denoted by $(kL)^{[n]}$. The following corollary was first obtained in [20] from Fujita’s approximation theorem (see [30] for further references). The proof given here combines theorem 1.4 with properties of zeroes of “random sections” [33].

Corollary 1.5. *If L is a big line bundle then*

$$\text{Vol}(L) = \lim_{k \rightarrow \infty} \frac{(kL)^{[n]}}{k^n}$$

The final two theorems concern the full Bergman kernel $K_k(x, y)$. First, the weak convergence of the squared point-wise norm of the $K_k(x, y)$ is obtained:

Theorem 1.6. *Let L be a line bundle and let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, E(k))$. Then*

$$k^{-n} |K_k(x, y)|_{k\phi}^2 \omega_n(x) \wedge \omega_n(y) \rightarrow \Delta \wedge \mu_\phi ,$$

as measures on $X \times X$, in the weak $*$ -topology, where Δ is the current of integration along the diagonal in $X \times X$.

Then a generalization of the Tian-Zelditch-Catlin expansion [37] for a globally positively curved line bundle is shown to hold for any (big) Hermitian line bundle L over a compact manifold X :

Theorem 1.7. *Let L be a line bundle and let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, E(k))$. Any interior point in $D \cap X(0) - \mathbb{B}_+(L)$ has a neighbourhood U where $K_k(x, y)e^{-k\phi(x)/2}e^{-k\phi(y)/2}$ (with x, y in U) admits an asymptotic expansion as*

$$(1.7) \quad k^n(\det(dd^c\phi)(x) + b_1(x, y)k^{-1} + b_2(x, y)k^{-2} + \dots)e^{k\phi(x, y)},$$

where b_i are global well-defined functions expressed as polynomials in the covariant derivatives of $dd^c\phi$ (and of the curvature of the metric ω) which can be obtained by the recursion given in [8].

Note that 1.6 in theorem 1.4 implies that $K_k(x, y)e^{-k\phi(x)/2}e^{-k\phi(y)/2} = |K_k(x, y)|_{k\phi}$ is exponentially small as soon as x or y is in the complement of D .

Remark 1.8. The assumption on ϕ may be relaxed to assuming that ϕ is in the class $\mathcal{C}^{1,1}$. For example, the proof of the regularity theorem 1.1 still goes through and the local Morse inequalities (lemma 4.1) still apply (almost everywhere on X). Moreover, all results remain true (with essentially the same proofs) if X is only assumed to be Moishezon, i.e. bimermorphically equivalent to a projective manifold (or equivalently, if L carries some big line bundle). However, in the remaining cases one would have to prove that $1_D \det(dd^c\phi) = 0$ almost everywhere on X . For example, if (X, ω) is a Kähler manifold then this would follow from the following conjecture:

Conjecture 1.9. *Let ω' be a smooth form cohomologous to the Kähler form ω . Then the global extremal function $V_{X, \omega'}$ associated to (X, ω') [24] (locally expressed as $\phi'_e - \phi'$, where $\omega' = dd^c\phi'$) is in the class $\mathcal{C}^{1,1}$.*

1.2. Further comparison with previous results. The present paper can be seen as a global geometric version of the situation recently studied in [5], where the role of the Hilbert space $\mathcal{H}(X, L^k)$ was played by the space of all polynomials in $\mathbb{C}^n$ of total degree less than k , equipped with a weighted norm (compare section 5). For references to previous works concerning the complex plane see [5]. The proof of the $\mathcal{C}^{1,1}$ -regularity of the equilibrium metric (on the complement of the augmented base locus of L) is partly modeled on the proof of Bedford-Taylor [2, 29] for $\mathcal{C}^{1,1}$ -regularity of the solution of the Dirichlet problem (with smooth boundary data) for the complex Monge-Ampere equation in the unit-ball in $\mathbb{C}^n$. The result should also be compared to various $\mathcal{C}^{1,1}$ -results for boundary value problems for complex Monge-Ampere equations on manifolds with boundary [13, 14], intimately related to the study of the geometry of the space of Kähler metrics on a Kähler manifold (see also

[31, 9] for other relations to Bergman kernels in the latter context). However, the present situation rather corresponds to a *free* boundary value problem (compare remark 3.10).

Further references and comments on the relation to the study of random polynomials (and holomorphic sections), random eigenvalues of normal matrices and various diffusion-controlled growth processes studied in the physics literature can be found in [5].

1.3. Further generalizations. In a sequel to this paper [7] *subspace* and *restricted versions* of the results in this paper will be obtained. The subspace version is a generalization of the case when the Hilbert space $\mathcal{H}(X, L^k)$ is replaced by the subspace of all sections vanishing to high order along a fixed divisor in X considered in the preprint [6]. Fixing a singular metric ϕ_s on L (with analytic singularities) the Hilbert space $\mathcal{H}(X, L^k)$ is replaced with the subspace of all global holomorphic sections of the twisted multiplier ideal sheaf $\mathcal{O}(L^k \otimes \mathcal{I}(k\phi_s))$ (i.e. the space of all sections f_k such that the point-wise norm $|f_k|^2 e^{-k\phi_s}$ is locally integrable) equipped with the Hilbert subspace norm in $\mathcal{H}(X, L^k)$ (i.e. the norm induced by the smooth metric ϕ). Similarly, the equilibrium metric ϕ_e is replaced by the metric obtained by further demanding that $\tilde{\phi} \leq \phi_s + C$ (i.e. that $\tilde{\phi}$ be more singular than ϕ_s) in the definition 3.1 of ϕ_e . As a special case new proofs of the results of Shiffman-Zelditch [35] about Hilbert spaces of polynomials with coefficients in a scaled Newton polytope are obtained.

The *restricted versions* are obtained by fixing an m -dimension complex submanifold V of X and replacing $\mathcal{H}(X, L^k)$ with the restricted space $\mathcal{H}(X, L^k)_V$ equipped with the “restricted norm” obtained by integrating sections over V . Similarly, the equilibrium metric ϕ_e is replaced by the metric defined on the restricted line bundle L_V by only demanding that $\tilde{\phi} \leq \phi$ on V in the definition 3.1. The corresponding Bergman kernel asymptotics can then be seen as local metrized versions of the very recent result in [21] concerning a generalized Fujita approximation theorem for the restricted volume (i.e. the asymptotic normalized dimension of $\mathcal{H}(X, L^k)_V$).

Acknowledgement 1.10. It is a pleasure to thank Jean-Pierre Demailly and Sebastian Boucksom for several illuminating discussions. Also thanks to Frederic Faure for his interest in this work and for drawing the pictures. This work was supported by a Marie Curie Intra European Fellowship.

1.4. General notation¹. Let (L, ϕ) be an Hermitian holomorphic line bundle over a compact complex manifold X . The fixed Hermitian fiber metric on L will be denoted by ϕ . In practice, ϕ is considered as a collection of local *smooth* functions. Namely, let s^U be a local holomorphic trivializing section of L over an open set U then locally, $|s^U(z)|_\phi^2 =: e^{-\phi^U(z)}$, where ϕ^U is in the class $\mathcal{C}^2$, i.e. it has continuous derivatives of order

¹general references for this section are the books [23, 16].

two. If α_k is a holomorphic section with values in L^k , then over U it may be locally written as $\alpha_k = f_k^U \cdot (s^U)^{\otimes k}$, where f_k^U is a local holomorphic function. In order to simplify the notation we will usually omit the dependence on the set U . The point-wise norm of α_k may then be locally expressed as

$$(1.8) \quad |\alpha_k|_{k\phi}^2 = |f_k|^2 e^{-k\phi}.$$

The canonical curvature two-form of L is the global form on X , locally expressed as $\partial\bar{\partial}\phi$ and the normalized curvature form $i\partial\bar{\partial}\phi/2\pi = dd^c\phi$ (where $d^c := i(-\partial + \bar{\partial})/4\pi$) represents the first Chern class $c_1(L)$ of L in the second real de Rham cohomology group of X . The curvature form of a smooth metric is said to be *positive* at the point x if the local Hermitian matrix $(\frac{\partial^2\phi}{\partial z_i \partial \bar{z}_j})$ is positive definite at the point x (i.e. $dd^c\phi_x > 0$). This means that the curvature is positive when $\phi(z)$ is strictly *plurisubharmonic* i.e. strictly subharmonic along local complex lines. We let

$$X(0) := \{x \in X : dd^c\phi_x > 0\}$$

More generally, a metric ϕ' on L is called (possibly) *singular* if $|\phi'|$ is locally integrable. Then the curvature is well-defined as a $(1,1)$ -current on X . The curvature current of a singular metric is called *positive* if ϕ' may be locally represented by a plurisubharmonic function (in particular, ϕ' takes values in $[-\infty, \infty[$ and is upper semi-continuous (u.s.c)). In particular, any section α_k as above induces such a singular metric on L , locally represented by $\phi' = \frac{1}{k} \ln |f_k|^2$. If Y is a complex manifold we will denote by $PSH(Y)$ and $SPSH(Y)$ the space of all plurisubharmonic and strictly plurisubharmonic functions, respectively.

Fix another line bundle F with a smooth metric ϕ_F and consider the following sequence of Hermitian holomorphic line bundles:

$$E(k) = (L^k \otimes F, k\phi + \phi_F)$$

Fixing an Hermitian metric two-form ω on X (with associated volume form ω_n) the Hilbert space $\mathcal{H}(X, E(k))$ is defined as the space $H^0(X, E(k))$ with the norm

$$(1.9) \quad \|\alpha_k\|_{k\phi}^2 (= \int_X |f_k|^2 e^{-(k\phi(z) + \phi_F)} \omega_n),$$

using a suggestive notation in the last equality (compare formula 1.8).

2. PRELIMINARIES: POSITIVITY AND BASE LOCI

Let L be a holomorphic line bundle over a compact projective Hermitian manifold (X, ω) .

2.1. Positivity for line bundles and singular metrics. The following notions of positivity will be used in the sequel [19]:

Definition 2.1. The line bundle L is said to be

(i) *pseudo-effective* if it admits a metric ϕ' with positive curvature current:

$$dd^c \phi' \geq 0$$

(ii) *big* if it admits a metric ϕ' with *strictly positive curvature current*:

$$(2.1) \quad dd^c \phi' \geq \epsilon \omega$$

(iii) *ample* if it admits a *smooth* metric ϕ' with *strictly positive curvature form*:

$$dd^c \phi'_x > 0$$

for all x in X .

2.2. Base loci. For each fixed k , the base locus of the line bundle $E(k)$ is defined as

$$\text{Bs}(|E(k)|) = \bigcap_{f_k \in H^0(X, E(k))} \{f_k = 0\}$$

(or as the corresponding ideal). *The stable base locus* $\mathbb{B}(L)$ of a line bundle L is defined [30] as the following analytic subvariety of X :

$$\mathbb{B}(L) := \bigcap_{k > 0} \text{Bs}(|kL|) = \bigcap_{f_k \in H^0(X, L^k), k \in \mathbb{N}} \{f_k = 0\}$$

In other words, a point x is in $\mathbb{B}(L)$ precisely when there is some section f_k in $H^0(X, L^k)$, for some k , which is non-vanishing at x . Moreover, the *augmented base locus* $\mathbb{B}_+(L)$ is defined in the following way [30]. Fix an ample line bundle on X . Then

$$\mathbb{B}_+(L) := \mathbb{B}(L - \epsilon A)$$

for any sufficiently small rational number ϵ (suitable interpreted using additive notation for tensor products). We will have great use for the following equivalent analytic definition of $\mathbb{B}_+(L)$ introduced in [11]) (and there called the *non-Kähler locus*)

$$(2.2) \quad X - \mathbb{B}_+(L) = \{x \in X : \exists \text{ big metric } \phi' \text{ on } L, \text{ smooth at } x\},$$

in the sense that ϕ' satisfies 2.1 and is smooth on some neighbourhood of the point x .² In fact, the equivalence of the definitions is a direct consequence of theorem 2.4 below. It amounts to showing that $\mathbb{B}_+(L)$ is the intersection of all effective divisors $E_k (= \{f_k = 0\})$ appearing in a “Kodaira decomposition”

$$(2.3) \quad L^k = A \otimes [E_k],$$

²the condition that ϕ' be smooth at x may be replaced by having 0 Lelong number at x .

for some positive natural number k . The point is that give any such decomposition

$$(2.4) \quad \phi_+ := \frac{1}{k} \ln |f_k|^2 + \phi_A$$

(where ϕ_A is a fixed smooth metric with positive curvature on A) is a metric on L with *strictly* positive curvature current such that ϕ_+ is *smooth* on $X - E$. The reason for calling $\mathbb{B}_+$ the augmented base locus is that

$$\mathbb{B}(L) \subseteq \mathbb{B}_+(L), \quad \bigcap_{k>0} \text{Bs}(|E(k)|) \subseteq \mathbb{B}_+(L)$$

Remark 2.2. It is well-known [11] that a line bundle L is ample precisely when $\mathbb{B}_+(L) = \emptyset$ and L is big precisely when $\mathbb{B}_+(L) \neq X$. In particular, L is non-ample and big precisely when $\mathbb{B}_+(L)$ is a non-empty analytic subvariety of X .

2.3. Extension of sections. Let $(L', \phi_{L'})$ and $(F', \phi_{F'})$ be line bundles with singular metrics such that

$$(2.5) \quad dd^c \phi_{L'} \geq \omega/C \text{ and } dd^c \phi_{F'} \geq -C\omega$$

for some positive number C .³ Recall the following celebrated theorem about L^2 -estimates for the $\bar{\partial}$ -equation, which is the basic analytical tool in the theory [16].

Theorem 2.3. (*Kodaira-Hörmander-Demailly*). *Let $\phi_{L'}$ and $\phi_{F'}$ be (singular) metrics as in 2.5. Take k' sufficiently large. Then for any $\bar{\partial}$ -closed $(0, 1)$ -form g with values in $L'^{k'} \otimes F'$ such that*

$$\|g\|_{k'\phi_{L'}+\phi_{F'}}^2 < \infty,$$

there is a section u with values in $L'^{k'} \otimes F'$ such that

$$\bar{\partial}u = g \text{ and } \|u\|_{k'\phi_{L'}+\phi_{F'}}^2 \leq \frac{C}{k'} \|g\|_{k'\phi_{L'}+\phi_{F'}}^2,$$

where the constant C is independent of k' and g .

The following extension theorem is a well-known asymptotic version of the Ohsawa-Takegoshi theorem [19]:

Theorem 2.4. *Let $\phi_{L'}$ and $\phi_{F'}$ be (singular) metrics as in 2.5. Fix a point x in X and take k' sufficiently large. Then any element in $(L'^{k'} \otimes F')_x$ such that*

$$|\alpha_{k'}|_{k'\phi_{L'}+\phi_{F'}}^2(x) < \infty$$

extends to a section in $H^0(X, L'^{k'} \otimes F')$ such that

$$(2.6) \quad \|\alpha_{k'}\|_{k'\phi_{L'}+\phi_{F'}}^2 \leq C |\alpha_{k'}|_{k'\phi_{L'}+\phi_{F'}}^2(x)$$

where the constant C is independent of k' and $\alpha_{k'}(x)$.

³ $(F', \phi_{F'})$ could also be replaced by a smooth Hermitian holomorphic vector bundle.

The previous theorem will be used to extend sections from $X - \mathbb{B}_+(L)$. The point is that given any (reasonable) metric $k\phi_L + \phi_F$ on $L^k \otimes F$ the following simple lemma provides a “strictly positively curved perturbation” ψ_k to which theorem 2.3 and 2.4 apply:

Lemma 2.5. *Let ϕ_L be a singular metric on L with positive curvature current and let ϕ_F be a smooth metric on the line bundle F . For any given point x_0 in $X - \mathbb{B}_+(L)$ and k sufficiently large there is a singular metric ψ_k on $L^k \otimes F$ such that $\psi_k = k'\phi_{L'} + \phi_{F'}$ with $\phi_{L'}$ and $\phi_{F'}$ as in formula 2.5 and*

$$(2.7) \quad \sup_{U(x_0)} |(k\phi_L + \phi_F) - \psi_k| \leq C_{x_0}$$

for some neighbourhood $U(x_0)$ of x_0 . Moreover, if ϕ_L is a metric with minimal singularities (see remark 3.5), for example the equilibrium metric ϕ_e (definition 3.1), then it may further be assumed that

$$(2.8) \quad \psi_k \leq (k\phi_L + \phi_F)$$

on all of X .

Proof. By the definition of $X - \mathbb{B}_+(L')$ there is a metric ϕ_+ on L , smooth in some $U(x_0)$ with *strictly* positive curvature current on X . Let $(F', \phi_{F'}) = (L^{k-k'} \otimes F, (k - k')\phi_L + \phi_F)$ and $(L', \phi') = (L, \phi_+)$, for k' fixed. Then $L^k \otimes F = L^{k'} \otimes F'$ gets an induced metric

$$(2.9) \quad \psi_k = (k - k')\phi_L + \phi_{F'} + k'\phi_+$$

of the required form for $k'(\leq k)$ sufficiently large, satisfying 2.7. Finally, if ϕ_L is a metric with minimal singularities we may assume that $\phi_+ \leq \phi_L$ after subtracting a sufficiently large constant from ϕ_+ . Then 2.8 clearly holds. $\square$

3. EQUILIBRIUM MEASURES FOR LINE BUNDLES

Let L be a line bundle over a compact complex manifold X . Given a smooth metric ϕ on L the corresponding “equilibrium metric” ϕ_e is defined as the envelope

$$(3.1) \quad \phi_e(x) = \sup \left\{ \tilde{\phi}(x) : \tilde{\phi} \in \mathcal{L}_{(X,L)}, \tilde{\phi} \leq \phi \text{ on } X \right\}.$$

where $\mathcal{L}_{(X,L)}$ is the class consisting of all (possibly singular) metrics on L with positive curvature current. Then ϕ_e is also in the class $\mathcal{L}_{(X,L)}$ (proposition 3.3 below). The Monge-Ampere measure $(dd^c \phi_e)^n / n!$ is well-defined on the open set

$$U(L) := \{x : \phi_e \text{ is bounded on } U(x)\},$$

where $U(x)$ is some neighbourhood of x (see [2, 28, 24] for the definition of the Monge-Ampere measure of a locally bounded metric or plurisubharmonic function). *The equilibrium measure* (associated to the metric ϕ) is now defined as

$$(3.2) \quad \mu_\phi := 1_{U(L)}(dd^c\phi_e)^n/n!$$

and is hence a positive measure on X . Consider the following set

$$(3.3) \quad D := \{\phi_e = \phi\} \subset X,$$

which is closed by (i) in the following proposition.

Proposition 3.1. *The following holds*

- (i) ϕ_e is in the class $\mathcal{L}_{(X,L)}$.
- (ii) $1_{U(L)}(dd^c\phi_e)^n/n! = 0$ on $X - D$.
- (iii) $D \subset \{x : dd^c\phi_x \geq 0\}$.

Proof. (i) is obtained by combining theorem 5.2 (2) and proposition 5.6 in [24].

The property (ii) is proved precisely as in the local theory in $\mathbb{C}^n$ (compare lemma 2.3 in the appendix of [32]). Indeed, it is enough to prove the vanishing on any small ball in $X - D$. For an alternative proof, using Bergman kernels, see the remark below.

To prove (iii) fix a point x where $dd^c\phi_x < 0$. Then there is a positive number ϵ and local coordinates z centered at x such that $(\frac{\partial^2\phi}{\partial\zeta\partial\bar\zeta})(\zeta, 0, \dots, 0) \leq \epsilon$ (with $z_1 = \zeta$) for ζ in the unit-disc Δ . Now take a candidate $\tilde{\phi}$ for the sup 3.1 and let $\psi_\epsilon(\zeta) := \tilde{\phi}(\zeta, 0, \dots, 0) - \phi(\zeta, 0, \dots, 0) - \epsilon|\zeta|^2$. Then $\psi_\epsilon(\zeta) \leq -\epsilon$ on $\partial\Delta$ and $\frac{\partial^2\psi_\epsilon}{\partial z_1\partial\bar{z}_1} \geq 0$ on Δ . Hence, the submean inequality for subharmonic functions (or the maximum principle) applied to ψ_ϵ gives $\tilde{\phi}(x) - \phi(x) = \psi_\epsilon(0) \leq -\epsilon$. Taking the sup over all candidates $\tilde{\phi}$ then gives $\phi_e(x) - \phi(x) \leq -\epsilon$ which proves the proposition. $\square$

Remark 3.2. As will be shown below ϕ_e is in the class $\mathcal{C}^1$ and its second derivatives exist almost everywhere and are locally bounded on $X - \mathbb{B}_+(L)$. Hence, one could also take $1_{X-\mathbb{B}_+(L)} \det(dd^c\phi_e)\omega_n$ as a somewhat more concrete definition of the equilibrium measure on X (which a posteriori anyway gives the same measure on X , according to theorem 3.4 below). It is interesting to see that the vanishing of $1_{X-\mathbb{B}_+(L)} \det(dd^c\phi_e)\omega_n$ on $X - D$ (corresponding to (ii) in the previous proposition) becomes a corollary of the proof of theorem 4.6 (which is independent of the proof of (ii) in the previous proposition).

Proposition 3.3. *The following properties of equilibrium metrics hold*

- (i) $(m\phi)_e = m\phi_e$
- (ii) Let ϕ_A be a metric on a line bundle A such that $dd^c\phi_A \geq 0$. Then

$$D_\phi \subseteq D_{\phi+\phi_A}$$

(iii) Assume that L is big and let ϕ_F be a smooth metric on a line bundle F . Then for any compact subset of $X - \mathbb{B}_+(L)$ there is a constant C such that

$$\frac{C}{m} - \phi_e \leq (\phi + \frac{1}{m}\phi_F)_e - \frac{1}{m}\phi_F \leq \phi_e + \frac{C}{m}$$

for all positive natural numbers m .

Proof. (i) is trivial. For (ii) note that $\phi_e + \phi_A$ is a contender for the sup in the definition of $(\phi + \phi_A)_e$. Hence, for x in D_ϕ we get

$$\phi(x) + \phi_A(x) = \phi_e(x) + \phi_A(x) \leq (\phi + \phi_A)_e(x).$$

This means that x is also in $D_{\phi+\phi_A}$, proving (ii).

To prove (iii) fix a compact set Ω in $X - \mathbb{B}_+(L)$ and a metric ϕ_+ on L with strictly positive curvature current, such that ϕ_+ is smooth Ω with $\phi_+ \leq \phi$ on X . Let us first prove one side of the inequality, i.e.

$$(3.4) \quad \phi_m := (\phi + \frac{1}{m}\phi_F)_e - \frac{1}{m}\phi_F \leq \phi_e + \frac{C}{m}$$

To this end first note that ϕ_m is a metric on L and there is clearly a constant C such that

$$\phi_m \leq \phi, \quad dd^c \phi_m \geq -\frac{C}{m} dd^c \phi_+.$$

Now let $\phi_{m,+} := (1 - \frac{C}{m})\phi_m + \frac{C}{m}\phi_+$, defining another metric on L . Note that

$$dd^c \phi_{m,+} \geq (1 - \frac{C}{m})(-\frac{C}{m} dd^c \phi_+) + \frac{C}{m} dd^c \phi_+ = (\frac{C}{m})^2 dd^c \phi_+ > 0$$

and since also $\phi_{m,+} \leq \phi$, the extremal definition of ϕ_e forces $\phi_{m,+} \leq \phi_e$. But since ϕ_+ is smooth on the compact set Ω this proves 3.4. The other side of the inequality is obtained from 3.4 applied to $\phi' = \phi + \frac{1}{m}\phi_F$ and $\phi'_F = -\phi_F$. $\square$

The next theorem gives the regularity properties of the equilibrium metric ϕ_e .

Theorem 3.4. *Suppose that L is a big line bundle and that the given metric ϕ on L is smooth (i.e. in the class $\mathcal{C}^2$). Then*

(a) ϕ_e is locally in the class $\mathcal{C}^{1,1}$ on $X - \mathbb{B}_+(L)$ i.e. ϕ_e is differentiable and all of its first partial derivatives are locally Lipschitz continuous there.

(b) The Monge-Ampere measure of ϕ_e on $X - \mathbb{B}_+(L)$ is absolutely continuous with respect to any given volume form and coincides with the corresponding $L_{loc}^\infty(n, n)$ -form obtained by a point-wise calculation:

$$(3.5) \quad (dd^c \phi_e)^n = \det(dd^c \phi_e) \omega_n$$

(c) the following identity holds almost everywhere on the set $D - \mathbb{B}_+(L)$, where $D = \{\phi_e = \phi\}$:

$$(3.6) \quad \det(dd^c \phi_e) = \det(dd^c \phi)$$

More precisely, it holds for all x in $D - \mathbb{B}_+(L) - G$, where G is the set defined in the proof of (c).

(d) Hence, the following identity between measures on X holds:

$$(3.7) \quad n! \mu_\phi = 1_{X - \mathbb{B}_+(L)} (dd^c \phi_e)^n = 1_D (dd^c \phi)^n = 1_{D \cap X(0)} (dd^c \phi)^n$$

Proof. (a), (b) and (c) will be proven in the subsequent section. To prove (d) first observe that the last equality in 3.7 follows immediately from proposition 3.1(iii). The second equality in 3.7 is obtained by combining (c) in the stated theorem with the vanishing in proposition 3.1(ii). Alternatively, the vanishing is obtained by combining the bound 4.13 applied to $\phi_e = \phi'_e$ with theorem 4.6, giving

$$1_{X-\mathbb{B}_+(L)}(dd^c\phi_e)^n = 1_D(dd^c\phi_e)^n$$

Finally, to obtain the vanishing of $(dd^c\phi_e)^n$ on $U(L) \cap \mathbb{B}_+(L)$ one can use the well-known local fact [28] that the Monge-Ampere measure of a locally bounded psh function integrates to zero over any pluripolar set (in particular over any local piece of $\mathbb{B}_+(L)$). $\square$

Remark 3.5. For a general line bundle L the equilibrium metric ϕ_e is an example of a metric with *minimal singularities* in the sense that for any other metric ϕ' in $\mathcal{L}_{(X,L)}$ there is a constant C such that

$$\phi' \leq \phi_e + C$$

on X (when such an inequality holds ϕ' is said to be *more singular* than ϕ_e) Such metrics play a key role in complex geometry [19].

3.1. The proof of $\mathcal{C}^{1,1}$ -regularity away from the augmented base locus. As in [5], where the manifold X was taken as $\mathbb{C}^n$, the proof is modeled on the proof of Bedford-Taylor [2, 29, 17] for $\mathcal{C}^{1,1}$ -regularity of the solution of the Dirichlet problem (with smooth boundary data) for the complex Monge-Ampere equation in the unit-ball in $\mathbb{C}^n$. However, as opposed to $\mathbb{C}^n$ and the unit-ball a generic compact Kähler manifold X has no global holomorphic vector fields. In order to circumvent this difficulty we will reduce the regularity problem on X to a problem on the manifold Y , where Y is the total space of the dual line bundle L^* , identifying the base X with its embedding as the zero-section in Y . To any given (possibly singular) metric ϕ on L we may associate the logarithm χ_ϕ of the “squared norm function” on Y , where locally

$$(3.8) \quad \chi_\phi(z, w) = \ln(|w|^2) + \phi(z),$$

in terms of local coordinates z_i on the base X and w along the fiber of L^* . In this way we obtain a bijection

$$(3.9) \quad \mathcal{L}_{(X,L)} \leftrightarrow \mathcal{L}_Y, \quad \phi \mapsto \chi_\phi$$

where $\mathcal{L}_Y$ is the class of all positively logarithmically 2-homogeneous plurisubharmonic functions on Y :

$$(3.10) \quad \mathcal{L}_Y := \{\chi \in PSH(Y) : \chi(\lambda \cdot) = \ln(|\lambda|^2) + \chi(\cdot)\},$$

using the natural multiplicative action of $\mathbb{C}^*$ on the fibers of Y over X . Now we define

$$(3.11) \quad \chi_e := \sup \{\chi \in \mathcal{L}_Y : \chi \leq \chi_\phi \text{ on } Y\}.$$

Then clearly, χ_e corresponds to the equilibrium metric ϕ_e under the bijection 3.9.

In the following we will denote by π the projection from Y onto X and by j the natural embedding of X in Y . We will fix a point y_0 in $Y - (j(X) \cup \pi^{-1}(\mathbb{B}_+(L)))$. Then there is a divisor E (appearing in a Kodaira decomposition as in formula 2.3) such that y_0 is in $Y - (j(X) \cup \pi^{-1}(E))$. Since clearly $(k\phi)_e = k\phi_e$ we may (since we are only interested in the regularity of ϕ_e) without loss of generality assume that k appearing in formula 2.3 is equal to one. Moreover, we fix an associated metric ϕ_+ (as in formula 2.4) that we write as

$$\phi_+ := \phi_A + \ln |e|^2,$$

using the suggestive notation e for the defining section of E , and $e(z)$ for any local representative. We may assume that $\phi_+ \leq \phi_e$. Later we will also assume the normalization $\chi_{\phi_e}(y_0) = 0$.

Existence of vector fields. The next lemma provides the vector fields needed in the modification of the approach of Bedford-Taylor.

Lemma 3.6. *Assume that the line bundle L is big. For any given point y_0 in $Y - [j(X) \cup \pi^{-1}(E)]$ there are global holomorphic vector fields $V_1, \dots, V_{n+1}$ (i.e. elements of $H^0(Y, TY)$) such that their restriction to y_0 span the tangent space TY_{y_0} . Moreover, given any positive integer m the vector fields may be chosen to satisfy*

$$(3.12) \quad (i) |V_i| \leq C_m(|w|)^m, \quad (ii) |V_i| \leq C_m(|e(z)|)^m$$

locally on the set $\{\chi_{\phi_+} \leq 1\}$ in Y (in the following we will fix some V_i corresponding to $m = 2$).

Proof. First note that Y may be compactified by the following fiber-wise projectivized vector bundle:

$$\widehat{Y} := \mathbb{P}(L^* \oplus \underline{\mathbb{C}}),$$

where $\underline{\mathbb{C}}$ denotes the trivial line bundle over X . Denote by $\mathcal{O}(1)$ the line bundle over $\widehat{Y}$ whose restriction to each fiber (i.e. a one-dimensional complex space $\mathbb{P}^1$) is the induced hyperplane line bundle. Next, we equip the line bundle

$$\widehat{L} := (\pi^*(L^{k_0}) \otimes \mathcal{O}(1))$$

over $\widehat{Y}$, where π denotes the natural projection from $\widehat{Y}$ to X , with a metric $\widehat{\phi}$ defined in the following way. First fix a smooth metric ϕ_E on the line bundle $[E]$ over X . Then

$$\phi_{+,k_0} := (\phi_A - \frac{1}{\sqrt{k_0}}\phi_E) + (1 + \frac{1}{\sqrt{k_0}})\ln |e|^2$$

is a metric on L such that $dd^c\phi_+ \geq dd^c\phi_A/2$ for $k_0 \gg 1$. Hence,

$$\widehat{\phi} := \pi^*(k_0\phi_{+,k_0}) + \ln(1 + e^{\chi_\phi})$$

is a metric on $\widehat{L}$ over Y (extending to $\widehat{Y}$) with strictly positive curvature current on $\widehat{Y}$, if $k_0 \gg 1$. Now fix a point y_0 in $(Y - (j(X) \cup \pi^{-1}(E)))$.

We can apply (ii) in theorem 2.4 to the bundle $\widehat{L}^{k_1} \otimes T\widehat{Y}$ over $\widehat{Y}$ for k_1 sufficiently large. Restricting to Y in $\widehat{Y}$ then gives that $TY \otimes \pi^*(L)^{k_0 k_1}$ is globally generated on Y (since $\mathcal{O}(1)$ is trivial on Y). Finally, observe that

$$\pi^*(L) = (\pi^*(L^*))^{-1} = [X]^{-1},$$

where $[X]$ is the divisor in Y determined by the embedding of X as the base. Indeed, X is embedded as the zero-set of the tautological section of $\pi^*(L^*)$ over $Y (= L^*)$. Hence, the sections of $TY \otimes \pi^*(L)^{k_0}$ may be identified with sections in TY vanishing to order k_0 on X . This proves (i) in 3.12. Moreover, by construction the vector fields V_i satisfy

$$|V_i(z, w)| \leq C \left(|w|^{k_0} |e(z)|^{k_0(1 + \frac{1}{\sqrt{k_0}})} |w| \right)^{k_1}$$

on any fixed neighbourhood in Y over the divisor E in X . Choosing k_0 and k_1 sufficiently large then gives

$$|V_i(z, w)| \leq C_m (|w| |e(z)|)^{k_0 k_1 + 1} |e(z)|^m.$$

Since, the factor $|w| |e(z)|$ is bounded on the set $\{\chi_{\phi_A + \ln|e|^2} \leq 0\}$ this proves (ii) in 3.12. $\square$

Existence of the flow. For any given smooth vector field V on Y and compact subset K of Y , we denote by $\exp(tV)$ the corresponding flow which is well-defined for any “time” t in $[0, t_K]$, i.e. the family of smooth maps indexed by t such that

$$(3.13) \quad \frac{d}{dt} f(\exp(tV)(y)) = df[V]_{\exp(tV)(y)}$$

for any smooth function f and point y on Y . We will also use the notation $\exp(V) := \exp(1V)$.

Combining the previous lemma with the inverse function theorem gives local “exponential” holomorphic coordinates centered at y_0 , i.e a local biholomorphism

$$\mathbb{C}^{n+1} \rightarrow U(y), \quad \lambda \mapsto \exp((V(\lambda)(y_0), \quad V(\lambda) := \sum \lambda_i V_i)$$

We will write

$$f^\lambda = (\exp(V(\lambda)))^* f$$

for the induced additive action on functions f (where the flow is defined). Using that the vector fields V_i necessarily also span TY_{y_1} for y_1 close to y_0 it can be checked that in order to prove that a function f is locally Lipschitz continuous on a compact subset of Y it is enough to, for each fixed point y_0 , prove an estimate of the form

$$(3.14) \quad |f^\lambda(y_0) - f(y_0)| \leq C |\lambda|$$

for some constant C only depending on the function f . Since we will later take f to be equal to χ_{ϕ_e} we may also, by homogeneity, assume that $\chi_{\phi_e}(y_0) = 0$. In order to define the flow on a neighbourhood U of

the whole levelset $\{\chi_{\phi_e} = 0\}$ (which is non-compact unless ϕ_e is locally bounded) we will use a compactification argument:

Lemma 3.7. *There is a positive number t_0 such that the flow $\exp(V(\lambda)y)$ exists for any (λ, y) such that $|\lambda| \leq t_0$ and y is in $U := \{\chi_{\phi_+} \leq 1\}$. Moreover, if ϕ' is a fixed metric on L such that $\phi' - \delta \ln |e|^2$ is smooth, for some number δ , then there are constants C_α such that*

$$(3.15) \quad |\partial_{z,w}^\alpha (\chi_{\phi'}^\lambda - \chi_{\phi'})| \leq C_\alpha |\lambda|$$

on $U \cap \pi^{-1}(X - E)$ over any fixed z -coordinate ball in X , in terms of the real local derivatives of multi order α and total order less than two.

Proof. Denote by Y' the total space of the ample line bundle $L \otimes [E]^{-1}$ over X and denote by π' the corresponding projection onto X . Then $Y \cap \pi^{-1}(X - E)$ is biholomorphic to $Y' \cap \pi'^{-1}(X - E)$ under the map Φ which may be locally represented as

$$(3.16) \quad \Phi : (z, w) \mapsto (z', w') := (z, e(z)w).$$

Note that Φ maps the set $\{\chi_{\phi_+} \leq 1\}$ to the set $\{\chi_{\phi_A} \leq 1\}$. Given a vector field V on $Y \cap \pi^{-1}(X - E)$ denote by V' the vector field $\Phi_* V$ on $Y' \cap \pi'^{-1}(X - E)$. Now fix a point in E corresponding to $z = 0$ in some local coordinates (z, w) for Y . Then the following local bound holds on the set $\{\chi_{\phi_A} \leq 1\}$ in Y' over $X - E$:

$$(3.17) \quad |V'_i(z', w')| \leq C |e(z')|.$$

Indeed, writing $V_i(z, w) = v_{i,z}(z, w) \frac{\partial}{\partial z} + v_{i,w}(z, w) \frac{\partial}{\partial w}$ and similarly for $V'_i(z', w')$ gives

$$(3.18) \quad v'_{i,z'}(z', w') = v_{i,z}(z, w), \quad v'_{i,w'}(z', w') = v_{i,z}(z, w) \frac{\partial e(z)}{\partial z} \frac{1}{e(z)} w' + v_{i,w}(z, w) e(z)$$

Hence, choosing vector fields V_i corresponding to $m = 2$ in (ii) in lemma 3.6 ensures that 3.17 holds.

Now fix a vector λ and write $\lambda = t\sigma$, where $|\lambda| = t$. Let V be the vector field on Y defined by the relation $V(\lambda) = tV$. By the local bounds 3.17, V' extends to a holomorphic vector field on the set $\{\chi_{\phi_A} \leq 1\}$ in Y' such that V' vanishes identically on $\pi'^{-1}(E)$. Since, $\{y' : \chi_{\phi_A}(y') \leq 1\}$ is compact in Y' the flow $\exp(tV')(y')$ is well-defined for $|t| \leq t_0$. Moreover, since V' vanishes identically on $\pi'^{-1}(E)$ the set $\pi'^{-1}(X - E)$ is invariant under the flow. By the isomorphism Φ in 3.16 this proves the existence of the flow stated in the lemma under the assumption that $y \in \pi^{-1}(X - E)$. But by the local bound (ii) in lemma 3.6 the flow does extend holomorphically over $\pi^{-1}(X - E)$.

To prove 3.15, note that since $\chi_{\phi'}$ is smooth over $X - E$ the defining property 3.13 of the flow gives

$$(3.19) \quad \chi_{\phi'}^\lambda - \chi_{\phi'} = \int_0^1 d\chi_{\phi'}[V(\lambda)]_{\exp(tV(\lambda)y)} dt$$

Hence, since $V(\lambda) := |\lambda| (\sum \frac{\lambda_i}{|\lambda|} V_i)$, the constant C_0 in 3.15 may be taken to be

$$C_0 = \sup_{y \in U \cap \pi^{-1}(X-E)} |d\chi_{\phi'}[V_i]_y| < \infty$$

To see that C_0 is finite it is, by the compactness of X , enough to prove the bound over any z -coordinate ball in X . First consider local coordinates (z, w) on Y where z is centered at a point in $X - E$. Then

$$(3.20) \quad |d\chi_{\phi'}[V_i]| = 2 \left| \frac{1}{w} dw[V_i] + \frac{\partial \phi'(z)}{\partial z} dz[V_i] \right| \leq C$$

using the bound (i) in lemma 3.6 for the first term in the right hand side above and the assumption that ϕ' is smooth on $X - E$ for the second term. Finally, consider the case when z is centered at a point in E . Then

$$\left| \frac{\partial \phi'(z)}{\partial z} \right| \leq C + \delta \left| \frac{1}{e(z)} \frac{\partial e(z)}{\partial z} \right| \leq C + C' \left| \frac{1}{e(z)} \right|.$$

Hence, the bound 3.20 still holds, using that $m(= 2) \geq 1$ in the local bound (ii) on V_i in lemma 3.6.

Finally note that the bounds for C_α when the total degree of α is two may be obtained in a completely similar maner, now using that $m = 2$ to handle the factors $\left| \frac{1}{e(z)} \right|^2$ and $\left| \frac{1}{w} \right|^2$. $\square$

Homogenization. To a given psh function $g(y)$ (defined on some disc subbundle of Y) we will associate the following S^1 -invariant psh function:

$$(3.21) \quad \widehat{g}(y) := \text{u.s.c.} \left(\sup_{\theta \in [0, 2\pi[} g(e^{i\theta} y) \right)$$

using the natural multiplicative action of $\mathbb{C}^*$ on the fibers of Y over X , where u.s.c. denotes the upper-semicontinuous regularization (using that the family $g(e^{i\theta} \cdot)$ of psh functions is locally bounded from above [28]).

The following simple lemma will allow us to “homogenize” in the normal direction of Y , as well:

Lemma 3.8. *Suppose that the function f is S^1 -invariant and psh on some disc subbundle U of Y containing the set $\{f \leq c\}$ in $(Y - \pi^{-1}(E))$, where E is an analytic variety in X . Moreover, assume that (i) f is strictly increasing along the fibers of Y over X and (ii) $f < c$ on the base X and $f > c$ on ∂U . Then there is a function $\tilde{f}$ in the class $\mathcal{L}_Y$ such that $\tilde{f} = f$ on the set $\{f = c\}$.*

Proof. It is enough to construct such a function $\tilde{f}$ on $Y - \pi^{-1}(E)$. Indeed, then $\tilde{f}$ is locally bounded from above close to $\pi^{-1}(E)$ and hence extends as a unique psh function to Y [28] (more generally E may be allowed to be locally pluripolar). Hence we may without loss of generality assume that E is empty in the following. First we will show that $f^{-1}(c)$ is an S^1 -subbundle of Y over X , i.e. the claim that the equation

$$f(\sigma) = c$$

has a unique solution on $\pi^{-1}(x)$, modulo the action of S^1 , for each fixed point x in X . To this end we identify the restriction of f to $\pi^{-1}(x)$ with a convex function $g(v)$ of $v = \ln |w|^2$ (using that f is S^1 -invariant and psh). In particular, g is continuous. By the assumption (i) the equation $g(v_0) = c$ has at most one solution. Moreover, by (ii) and the fact that g is continuous the solution v_0 does exist. This proves the claim above. Now define

$$\tilde{f}(r\sigma) := \ln(r^2) + f(\sigma).$$

To see that $\tilde{f}$ is psh, we may, since the problem is local, assume that X is a ball in $\mathbb{C}_z^n$. Write $\tilde{f}(z, w) = \tilde{\phi}(z) + \ln |w|^2$ and note that

$$\Omega = U \cap \{\tilde{\phi}(z) + \ln |w|^2 \leq c\} = U \cap \{f \leq c\}$$

using that f is strictly increasing along the fibers. Since f is psh it follows that $\partial\Omega$ is pseudoconvex. A classical result of Bremermann [12] for such Hartogs domains Ω now implies that $\tilde{\phi}(z)$ and hence $\tilde{f}(z, w)$ is psh.⁴ $\square$

Lemma 3.9. *For each λ (with sufficiently small norm) the function $f = \widehat{\chi_{\phi_e}^\lambda}$ satisfies the assumptions in the previous lemma with $c = c_\lambda = \widehat{\chi_{\phi_e}^\lambda}(y_0)$ and $U = \{\chi_{\phi_+} \leq 1\}$.*

Proof. Given local coordinates on U , let

$$f_\lambda(z, w) := \chi_{\phi_e}^\lambda(z, w) - \ln |w|^2 = [\phi_e^\lambda(z, w) + ((\ln |w|^2)^\lambda - \ln |w|^2)].$$

Since $\phi_e^\lambda(z, w)$ is psh (a local version of) 3.15 in lemma 3.7) applied to $\phi' = \ln |w|^2$ gives

$$\frac{\partial^2 f_\lambda(z, w)}{\partial w \partial \bar{w}} \geq -C |\lambda|$$

Note that the constant C may be taken to be independent of the local coordinates as in the proof of 3.15 in lemma 3.7, since the base X is compact. Now write

$$\widehat{\chi_{\phi_e}^\lambda} = \widehat{g} - C |\lambda| |w|^2 + \ln |w|^2$$

with $g = f_\lambda + C |\lambda| |w|^2$.⁵ By the maximum principle $\widehat{g}$ in the definition 3.21 is always increasing in $v := \ln |w|^2$. Hence the (right) derivative of the convex function $\widehat{\chi_{\phi_e}^\lambda}$ with respect to v (with the variable z fixed) is positive (and almost equal to 1) for λ sufficiently small. This proves (i).

To prove (ii) first observe that

$$(3.22) \quad c_\lambda := \widehat{\chi_{\phi_e}^\lambda}(y_0) \leq 1/2$$

⁴alternatively one can, by local approximation, reduce to the case when $\tilde{f}$ is smooth and c regular. Then $dd^c \tilde{f} = (\tilde{f}/f) dd^c f \geq 0$ along $T^{1,0}(\partial\Omega)$ and hence, by homogeneity, $\tilde{f}$ is psh on Y .

⁵over a neighbourhood of E we use the local coordinates (z', w') so that the function $|w'|^2$ is uniformly bounded on U .

for all λ (with sufficiently small norm). Indeed, by upper semi-continuity $\limsup_{\lambda \rightarrow 0} \widehat{\chi_{\phi_e}^\lambda}(y_0) \leq \widehat{\chi_{\phi_e}}(y_0) = 0$. Hence, 3.22 holds (after possibly replacing the upper bound t_0 on $|\lambda|$ with a smaller number). Next, note that by the extremal definition of ϕ_e we have $\widehat{\chi_{\phi_e}^\lambda} \geq \chi_{\phi_+}^\lambda$ on Y . By 3.22 and the definition of U it is hence enough to prove that $\chi_{\phi_+}^\lambda \geq \chi_{\phi_+} - C|\lambda|$ on U . But this follows from 3.15 in lemma 3.7 applied to $\phi' = \phi_+$. $\square$

3.1.1. *Proof of (a)–(c) in theorem 3.4.* To prove (a) it is, by the bijection 3.9, equivalent to prove that χ_e (defined by 3.11) is locally $\mathcal{C}^{1,1}$ on $Y - [j(X) \cup \pi^{-1}(\mathbb{B}_+(L))]$.

Proof. Step1: $\chi_e(:=\chi_{\phi_e})$ is locally Lipschitz continuous on $Y - (j(X) \cup \pi^{-1}(\mathbb{B}_+(L)))$.

To see this fix a point y_0 as above. From the definition 3.21 of $(\widehat{\cdot})$ we have an upper bound

$$\chi_{\phi_e}^\lambda(y_0) \leq \widehat{\chi_{\phi_e}^\lambda}(y_0) = \widetilde{\widehat{\chi_{\phi_e}^\lambda}}(y_0),$$

where $\widetilde{\widehat{\chi_{\phi_e}^\lambda}}$ is the function in the class $\mathcal{L}_Y$, extending $\widehat{\chi_{\phi_e}^\lambda}$ from the level set

$$M_\lambda := \{y : \widehat{\chi_{\phi_e}^\lambda}(y) = \widehat{\chi_{\phi_e}^\lambda}(y_0)\} \subset U$$

obtained from lemma 3.9. Since, by definition, $\chi_{\phi_e}^\lambda \leq \chi_\phi^\lambda$ we have the following bound on the level set M_λ :

$$(3.23) \quad \widetilde{\widehat{\chi_{\phi_e}^\lambda}}(y) \leq \sup_{\theta \in [0, 2\pi]} \chi_\phi((\exp(V(\lambda))(e^{i\theta}y)) \leq \sup_{\theta \in [0, 2\pi]} \chi_\phi(e^{i\theta}y) + C|\lambda|,$$

using that χ_ϕ is smooth in the last inequality so that 3.15 in lemma 3.7 can be applied. Since χ_ϕ is S^1 -invariant 3.23 gives that

$$(3.24) \quad \widetilde{\widehat{\chi_{\phi_e}^\lambda}} - C|\lambda| \leq \chi_\phi$$

on the level set M_λ and hence, by homogeneity, on all of Y . This shows that the function $\widetilde{\widehat{\chi_{\phi_e}^\lambda}} - C|\lambda|$ is a contender for the supremum in the definition 3.11 of χ_e and hence bounded by χ_e . All in all we get that

$$\chi_{\phi_e}^\lambda(y_0) \leq \widetilde{\widehat{\chi_{\phi_e}^\lambda}}(y_0) \leq \chi_{\phi_e}(y_0) + C|\lambda|.$$

The other side of the inequality 3.14 for $f = \chi_{\phi_e}^\lambda$ is obtained after replacing λ by $-\lambda$.

Step2: $d\chi_e$ exists and is locally Lipschitz continuous on $Y - X$.

Following the exposition in [17] of the approach of Bedford-Taylor it is enough to prove the following inequality:

$$(3.25) \quad (\chi_{\phi_e}^{-\lambda}(y_0) + \chi_{\phi_e}^\lambda(y_0))/2 - \chi_{\phi_e}(y_0) \leq C|\lambda|^2,$$

where the constant only depends on the second derivatives of χ_ϕ . Indeed, given this inequality (combined with the fact that χ_{ϕ_e} is psh) a Taylor

expansion of degree 2 gives the following bound close to y_0 for a local smooth approximation χ_ϵ of χ_{ϕ_e} :

$$|D^2\chi_\epsilon| \leq C$$

where $\chi_\epsilon := \chi_{\phi_e} * u_\epsilon$, using a local regularizing kernel u_ϵ and where $D^2\chi_\epsilon$ denotes the real local Hessian matrix of χ_{ϕ_e} . Letting ϵ tend to 0 then proves Step 2. Finally, to see that the inequality 3.25 holds we apply the argument in Step 1 after replacing $\chi_{\phi_e}^\lambda$ by the psh function

$$g(y) := (\chi_{\phi_e}^\lambda(y) + \chi_{\phi_e}^{-\lambda}(y))/2$$

to get the following bound on the level set $\{y : \widehat{g}(y) = \widehat{g}(y_0)\}$:

$$g(y) \leq \widetilde{g}(y) \leq \sup_{\theta \in [0, 2\pi]} (\chi_\phi((\exp(V(\lambda))(e^{i\theta}y)) + \chi_\phi(\exp(V(-\lambda))(e^{i\theta}y))/2$$

Next, observe that for each fixed θ the function $\chi_\phi(e^{i\theta}y)$ is in the class $\mathcal{C}^2$. Hence, a Taylor expansion of degree 2 in the right hand side of formula 3.19 gives

$$\widetilde{g}(y) \leq \sup_{\theta \in [0, 2\pi]} ((\chi_\phi(e^{i\theta}y)) + C|\lambda|^2) = \chi_\phi(y) + C|\lambda|^2$$

where the constant C may be taken as a constant times $\sup_U |\langle D^2\chi_\phi V, V \rangle_\eta|$ in terms of some fixed metric η on TY . This shows that $\widetilde{g} - C|\lambda|^2$ is a contender for the supremum in the definition 3.11 of χ_e and hence bounded by χ_e . All in all we obtain that

$$g(y_0) \leq \chi_\phi(y_0) + C|\lambda|^2,$$

which proves the inequality 3.25, finishing the proof of Step2.

(c) To see that 3.6 holds, it is enough to prove that locally

$$(3.26) \quad \frac{\partial^2 \phi}{\partial x_i \partial x_j}(\phi_e - \phi) = 0$$

almost everywhere on $D = \{\phi_e = \phi\}$, where x_i is a real coordinate on $\mathbb{R}^{2n} (= \mathbb{C}^n)$. To this end let $\psi := \phi_e - \phi$ and let $A := \{\psi = 0, d\psi \neq 0\}$. By (a) above ψ is a $\mathcal{C}^1$ -function and hence A is a real hypersurface of codimension 1 and in particular of measure zero (w.r.t. Lebesgue measure). Next, let $f := d\psi$ (considered as a local map on $\mathbb{R}^{2n}$) and let B_1 be the set where the derivative df (i.e the matrix $(df_1, \dots, df_{2n})$) does not exist. Since, by (a) f is a Lipschitz map it is well-known that B_1 also has measure zero. Finally, let B_2 be the set where $f = 0$, df exists, but $df \neq 0$. Now using a lemma in [27] (page 53) applied to the Lipschitz map f gives that B_2 too has measure zero. Finally, let $G := A \cup B_1 \cup B_2$. Then G has measure zero and 3.26 holds on $(X - \mathbb{B}_+(L)) - G$, proving (c). $\square$

Remark 3.10. Suppose that L is ample and fix a smooth metric ϕ_+ on L with positive curvature. Then $\omega_+ := dd^c \phi_+$ is a Kähler metric on X and the fixed metric ϕ on L may be written as $\phi = u + \phi_+$, where u

is a smooth *function* on X . Now the pair (u_e, M) where $u_e := \phi_e - \phi_+$ and M is the set $X - D$, may be interpreted as a “weak” solution to the following *free* boundary value problem of Monge-Ampere type⁶:

$$\begin{aligned} (dd^c u_e + \omega_+)^n &= 0 && \text{on } M \\ u_e &= u && \text{on } \partial M \\ du_e &= du \end{aligned}$$

The point is that, since the equations are overdetermined, the set M is itself part of the solution. In [25] the $\mathcal{C}^{1,1}$ –regularity of ϕ_e in the case when $X = \mathbb{C}$ (corresponding to the setup in [32]) was deduced from the regularity of a free boundary value problem.

4. BERGMAN KERNEL ASYMPTOTICS

Recall that $\mathcal{H}(X, E(k))$ denotes the Hilbert space obtained by equipping the vector space $H^0(X, E(k))$ with the norm 1.9. Let (ψ_i) be an orthonormal base for $\mathcal{H}(X, E(k))$. The *Bergman kernel* of the Hilbert space $\mathcal{H}(X, E(k))$ is the integral kernel of the orthogonal projection from the space of all smooth sections with values in $E(k)$ onto $\mathcal{H}(X, E(k))$. It may be represented by the holomorphic section

$$(4.1) \quad K_k(x, y) = \sum_i \psi_i(x) \otimes \overline{\psi_i(y)}.$$

of the pulled back line bundle $E(k) \boxtimes \overline{E(k)}$ over $X \times \overline{X}$. The restriction of K_k to the diagonal is a section of $E(k) \otimes \overline{E(k)}$ and we let $B_k(x) = |K_k(x, x)|_{k\phi + \phi_F} (= |K_k(x, x)| e^{-(k\phi(x) + \phi_F)})$ be its point wise norm:

$$(4.2) \quad B_k(x) = \sum_i |\psi_i(x)|_{k\phi + \phi_F}^2.$$

We will refer to $B_k(x)$ as the *Bergman function* of $\mathcal{H}(X, E(k))$. It has the following extremal property:

$$(4.3) \quad B_k(x) = \sup \left\{ |\alpha_k(x)|_{k\phi + \phi_F}^2 : \alpha_k \in \mathcal{H}(X, E(k)), \|\alpha_k\|_{k\phi + \phi_F}^2 \leq 1 \right\}$$

Moreover, integrating 4.2 shows that B_k is a “dimensional density” of the space $\mathcal{H}(X, L^k)$:

$$(4.4) \quad \int_X B_k \omega_n = \dim \mathcal{H}(X, E(k))$$

The following “local Morse inequality” estimates B_k point-wise from above for a general bundle:

Lemma 4.1. (Local Morse inequalities) *On any compact complex manifold the following upper bound holds:*

$$k^{-n} B_k \leq C_k 1_{X(0)} \det(dd^c \phi),$$

⁶since there is a priori no control on the regularity of the set M , it does not really make sense to write ∂M and the boundary condition should hence be interpreted in a suitable “weak” sense.

where the sequence C_k of positive numbers tends to one and $X(0)$ is the set where $dd^c\phi > 0$.

See [3] for the more general corresponding result for $\bar{\partial}$ -harmonic $(0, q)$ -forms with values in a high power of an Hermitian line bundle. The present case (i.e. $q = 0$) is a simple consequence of the mean-value property of holomorphic functions applied to a poly-disc Δ_k of radius $\ln k/\sqrt{k}$ centered at the origin in $\mathbb{C}^n$ (see the proof in [4]). In fact, the proof gives the following stronger local statement:

$$(4.5) \quad \limsup_k k^{-n} \left| f_k(z/\sqrt{k}) \right|^2 e^{-(k\phi + \phi_F)((z/\sqrt{k}))} / \|f_k\|_{k\phi + \phi_F, \Delta_k}^2 \leq 1_{X(0)}(0) \det(dd^c\phi),$$

where f_k is holomorphic function defined in a fixed neighbourhood of the origin in $\mathbb{C}^n$.

The local estimate in the previous lemma can be considerably sharpened on the complement of the globally defined set D (formula 3.3), as shown by the following lemma:

Lemma 4.2. *Let ϕ be a smooth metric on a holomorphic line bundle L over a compact manifold X .*

(i) if B_k denotes the Bergman function of the Hilbert space $\mathcal{H}(X, L^k)$, then the following inequality holds on all of X :

$$(4.6) \quad B_k k^{-n} \leq C_k e^{-k(\phi - \phi_e)}$$

where the sequence C_k of positive numbers tends to $\sup_X \det(dd^c\phi)$.

(ii) If L is big and B_k now denotes the Bergman function of the Hilbert space $\mathcal{H}(X, E(k))$, then the inequality 4.6 holds on any given compact subset of $X - \mathbb{B}_+(L)$ if C_k is replaced by a large constant C (depending on the compact set).

In particular, in both cases

$$(4.7) \quad \lim \int_{D^c} k^{-n} B_k \omega_n = 0$$

Proof. Let us first prove (i). By the extremal property 4.3 of B_k it is enough to prove the lemma with $B_k k^{-n}$ replaced by $|\alpha_k|_{k\phi}^2$, locally represented by $|f_k| e^{-k\phi}$, for any element α_k in $\mathcal{H}(X, L^k)$ with global norm equal to k^{-n} . The Morse inequalities in the previous lemma give that

$$|f_k|^2 e^{-k\phi} \leq C_k$$

with C_k as in the statement of the present lemma. Equivalently,

$$\frac{1}{k} \ln |f_k|^2 - \frac{1}{k} C_k \leq \phi$$

Hence, the singular metric on L determined by $\frac{1}{k} \ln |f_k|^2 - \frac{1}{k} C_k$ is a candidate for the sup in the definition 3.1 of ϕ_e and is hence bounded by ϕ_e . Thus,

$$B_k k^{-n} = |f_k|^2 e^{-k\phi} \leq C_k e^{k\phi_e} e^{-k\phi}.$$

The proof of (ii) is completely analogous if one takes into account that $\ln |f_k|^2$ is now a metric on $E(k) = L^k \otimes F$ and uses (ii) in proposition 3.3. Finally, the vanishing 4.7 follows from the dominated convergence theorem (using that the sequence $B_k k^{-n}$ is, by lemma 4.1, uniformly bounded on X), since the right hand side in the previous inequality tends point-wise to zero precisely on the complement of D . $\square$

Remark 4.3. When L is ample the supremum, in the definition of C_k , in the previous lemma can be taken over the support of $(dd^c \phi_e)^n$, using a max/comparison principle.

The following lemma yields a *lower* bound on the Bergman function.

Lemma 4.4. *Let L be a big line bundle, then the following lower bound holds at almost any point x in $D \cap X(0)$:*

$$(4.8) \quad \liminf_k k^{-n} B_k \omega_n \geq (dd^c \phi)^n / n!$$

Proof. Step1: construction of a smooth extremal σ_k . Fix a point x_0 in $D \cap X(0) - \mathbb{B}_+(L) - G$, where G is the set of measure zero appearing in the proof of (c) in the regularity theorem 3.4. First note that there is a *smooth* section σ_k with values in $E(k)$ such that

$$(4.9) \quad (i) \lim_{k \rightarrow \infty} \frac{|\sigma_k|_{k\phi}^2(x_0)}{k^n \|\sigma_k\|_{k\phi+\phi_F}^2} \omega_n = (dd^c \phi)_{x_0}^n, \quad (ii) \|\bar{\partial} \sigma_k\|_{k\phi_e+\phi_F}^2 \leq C e^{-k/C}$$

To see this take trivializations of L and F and local holomorphic coordinates z_i centered at x_0 (and orthonormal at x_0) such that $\phi_F(0) = 0$ and

$$(4.10) \quad \phi(z) = (\sum_{i=1}^n \lambda_i |z_i|^2 + O(|z|^3))$$

with λ_i the positive eigenvalues of $(dd^c \phi)_{x_0}$ w.r.t the metric ω [23]. Fix a smooth function χ which is constant when $|z| \leq \delta/2$ and supported where $|z| \leq \delta$; the number δ will be assumed to be sufficiently small later on. Now σ_k is simply obtained as the local section with values in L^k represented by the function χ close to x_0 and extended by zero to all of X . To see that (i) holds note that, using 4.10,

$$\lim_{k \rightarrow \infty} \frac{|\sigma_k|_{k\phi}^2(x_0)}{k^n \|\sigma_k\|_{k\phi+\phi_F}^2} = \lim_{k \rightarrow \infty} \frac{\chi(0)}{k^n \int_{|z| \leq k^{-1/2} \ln k} e^{-k \sum_{i=1}^n \lambda_i |z_i|^2} \chi(0) \omega_n(0)},$$

where $\omega_n(0)$ is the Euclidian volume form in $\mathbb{C}^n$ (since z_i are assumed to be orthonormal w.r.t ω at 0). Evaluating the latter Gaussian integral then gives the limit $(1/\pi)^n \lambda_1 \lambda_2 \cdots \lambda_n$, proving (i) in 4.9. To prove (ii) in 4.9, first note that

$$\|\bar{\partial} \sigma_k\|_{k\phi_e+\phi_F}^2 \leq C \int_{\delta/2 \leq |z| \leq \delta} e^{-k(\phi(z) + (\phi_e(z) - \phi(z)))} \omega_n(0),$$

as follows from the definition of χ . Hence, (ii) follows from the fact that

$$(4.11) \quad |z| \leq \delta \Rightarrow \phi(z) + (\phi_e(z) - \phi(z)) \geq \inf_i \lambda_i |z|^2 / 2$$

for δ sufficiently small, if x_0 is in the set $D \cap X(0) - \mathbb{B}_+(L) - G$. To prove 4.11 fix $z \neq 0$ and let $\psi(y) := \phi_e(y \frac{z}{|z|}) - \phi(y \frac{z}{|z|})$, where y is a non-negative number. By the regularity theorem 3.4 u is in the class $\mathcal{C}^1$. Hence, since x_0 is in D (where $\psi = 0$)

$$|\psi(y)| \leq \int_0^y \left| \frac{d\psi}{ds} \right| (s) ds.$$

Moreover, since x_0 is in $D - \mathbb{B}_+(L) - G$, we have $\frac{d\psi}{ds}(0) = 0$ and $\lim_{s \rightarrow 0} \frac{d\psi}{ds}(s)/s = 0$. In particular, for any given positive number ϵ we have that

$$|\psi(y)| \leq \int_0^y \epsilon s ds = \epsilon y^2 / 2$$

for $y \leq \delta$, if δ is sufficiently small. Combining the latter estimate with 4.10 then proves 4.11 and finishes the proof of (ii) in 4.9.

Step2: perturbation of σ_k to a holomorphic extremal α_k . Equip $E(k)$ with a “strictly positively curved modification” ψ_k of the metric $k\phi_e + \phi_F$ furnished by lemma 2.5. Let $g_k = \bar{\partial}\sigma_k$ and let α_k be the following holomorphic section

$$\alpha_k := \sigma_k - u_k,$$

where u_k is the solution of the $\bar{\partial}$ -equation in the Hörmander-Kodaira theorem 2.3 with $g_k = \bar{\partial}\sigma_k$. Hence,

$$\|u_k\|_{\psi_k} \leq C \|g_k\|_{\psi_k}$$

Next, applying 2.7 to the right hand side above (using that g_k is supported on a small neighborhood of $x_0 \in X - \mathbb{B}_+(L)$ and then 2.8 to the left hand side above gives

$$\|u_k\|_{k\phi_e + \phi_F} \leq C \|g_k\|_{k\phi_e + \phi_F}$$

Finally, using that $\phi_e \leq \phi$ on all of X in the left hand side above gives

$$(4.12) \quad \|u_k\|_{k\phi + \phi_F} \leq C \|g_k\|_{k\phi_e + \phi_F}$$

and then (ii) in 4.9 in the right hand side gives

$$(a) \|u_k\|_{k\phi + \phi_F} \leq C e^{-k/C}, \quad (b) |u_k|_{k\phi + \phi_F}^2(x) \leq C' k^n e^{-k/C'},$$

where (b) is a consequence of (a) and the local holomorphic Morse inequalities 4.5 applied to u_k at $z = 0$. Combining (a) and (b) with (i) in 4.9 then proves that (i) in 4.9 holds with σ_k replaced by the holomorphic section α_k . By the definition of B_k this finishes the proof of the lemma. $\square$

Remark 4.5. For *any* line bundle L over X and ϕ' a given (singular) metric on L with positive curvature form Boucksom showed [10] that

$$(4.13) \quad \liminf_k k^{-n} \dim H^0(X, E(k)) \geq \int_X ((dd^c \phi')_{ac})^n / n!,$$

where $(dd^c \phi')_{ac}$ denotes the absolutely continuous part of the the current $dd^c \phi'$. Boucksom used Bonavero's strong Morse inequalities for singular

metrics with analytic singularities. However, when L is big the lower bound 4.13 follows from a variant of the proof of the previous lemma. To see this one first approximates ϕ' with a sequence ϕ'_ϵ with *analytic* singularities (as in [10]) and replace ϕ with the metric

$$\phi'_{\epsilon,+} := \phi'_\epsilon(1 - \epsilon) + \epsilon\phi_+$$

with *strictly* positive curvature form (since ϕ_+ is of the form 2.1) and the Hilbert space $\mathcal{H}(X, E(k))$ with the Hilbert space $\mathcal{H}(X, E(k), \phi'_{\epsilon,+})$ whose norm is defined with respect to the norm induced by the singular metric $\phi'_{\epsilon,+}$.⁷ A variant of the proof of the previous lemma then gives a lower bound on the corresponding Bergman function with ϕ replaced by ϕ'_ϵ , for any point x in the complement of the singularity locus of ϕ'_ϵ . Finally, integrating over X and then letting ϵ tend to zero gives 4.13.

Theorem 4.6. *Let B_k be the Bergman function of the Hilbert space $\mathcal{H}(X, E(k))$. Then*

$$(4.14) \quad k^{-n} B_k(x) \rightarrow 1_{D \cap X(0)} \det(dd^c \phi)(x)$$

for almost any x in X , where $X(0)$ is the set where $dd^c \phi > 0$ and D is the set 3.3. Moreover, the following weak convergence of measures holds:

$$k^{-n} B_k \omega_n \rightarrow \mu_\phi,$$

where μ_ϕ is the equilibrium measure.

Proof. Case1: L is big

First observe that, by the exponential decay in lemma 4.2,

$$\lim_{k \rightarrow \infty} k^{-n} B_k(x) = 0, \quad x \in D^c$$

Next, the local Morse inequalities (lemma 4.1) give the upper bound in 4.14 on $k^{-n} B_k(x)$ for any x in D and lemma 4.4 gives, since L is assumed to be big, the lower bound for almost any x in D , finishing the proof of 4.14. Finally, the weak convergence follows from 4.14 combined with the uniform upper bound on $k^{-n} B_k(x)$ in lemma 4.1) (using the dominated convergence theorem on X).

Case2: L is pseudo-effective, but not big

First note that the dimension of $H^0(X, E(k))$ is of the order $o(k^n)$, i.e.

$$(4.15) \quad \lim_{k \rightarrow \infty} \int_X k^{-n} B_k \omega_n = 0.$$

To see this one argues by contradiction (compare proposition 6.6f in [19]) : if the dimension would be of the order k^n a standard argument gives that $H^0(X, E(k) \otimes A^{-1})$ has a section for k sufficiently large, if A is any fixed ample line bundle. But then

$$L^k = (A \otimes F^{-1}) \otimes E,$$

⁷as a vector space the Hilbert space $\mathcal{H}(X, E(k), \phi'_{\epsilon,+})$ consists of sections which have finite norm with respect to the singular metric $\phi'_\epsilon(1 - \epsilon) + \epsilon\phi_+$.

where E is an effective divisor. Since we may choose A so that $A \otimes F^{-1}$ is ample this means that L has a metric with *strictly* positive curvature form (compare formula 2.4), giving a contradiction.

Now given 4.15, Fatou's lemma forces

$$\lim_{k \rightarrow \infty} k^{-n} B_k \omega_n(x) = 0$$

for almost all x in X . Finally, to prove the vanishing of the positive measure $1_{D \cap X(0)} \det(dd^c \phi)(x)$ a.e. on X , equip the ample line bundle A with a metric ϕ_A with positive curvature form. Then proposition 3.3 (i) gives that

$$\int_{D_{p\phi} \cap X(0)} (dd^c p\phi)^n / n! \leq \int_{D_{p\phi + \phi_A} \cap X(0)} (dd^c (p\phi + \phi_A))^n / n!.$$

Using (ii) in proposition 3.3 and that $dd^c p\phi \leq dd^c (p\phi + \phi_A)$ then gives (4.16)

$$\int_{D \cap X(0)} (dd^c \phi)^n / n! \leq p^{-n} \int_{D_{p\phi + \phi_A} \cap X(0)} (dd^c (p\phi + \phi_A))^n / n! = p^{-n} \text{Vol}(pL + A),$$

where we have applied case 1 to the big line bundle $pL + A$ in the last step and used the definition 4.17 below of the volume $\text{Vol}(L')$. Now fix $\epsilon > 0$. By the ‘‘continuity’’ of the volume function (compare remark 4.10) the right hand side is bounded by

$$p^{-n} \text{Vol}(pL) + \epsilon = \epsilon.$$

Finally, letting p tend to infinity in 4.16 gives that $\int_{D \cap X(0)} (dd^c \phi)^n / n!$ must vanish, since ϵ was arbitrary.

Case3: L is not pseudo-effective (and hence not big)

In this case it follows directly from the definition that $\phi_e \equiv -\infty$ and hence the set D is empty. $\square$

Remark 4.7. As shown in the proof of lemma 4.4 the set G of measure zero where the point-wise convergence in the previous theorem may fail, can be expressed in terms of the ‘‘derivatives’’ (up to order two) of $\phi_e - \phi$. The proof of lemma 4.4 also shows that the L^1 -convergence for the corresponding Bergman functions for weighted polynomials obtained in [5] can be improved to convergence on the complement of G .

The volume of a line bundle is defined by the following formula [30]:

$$(4.17) \quad \text{Vol}(L) := \limsup_k k^{-n} \dim H^0(X, L^k)$$

Integrating the convergence of the Bergman kernel in 4.6 combined with the previous corollary gives the following version of Fujita's approximation theorem [22, 10]:

Corollary 4.8. *The volume of a big line bundle L is given by*

$$(4.18) \quad \text{Vol}(L) = \int_{X - \mathbb{B}_+(L)} (dd^c \phi_e)^n / n!$$

and $\text{Vol}(L) = 0$ precisely when L is not big.

Remark 4.9. Assume that L is big. Fujita's approximation theorem (as formulated in [30, 20]) may be stated as

$$(4.19) \quad \text{Vol}(L) = \sup_A A^n$$

where the supremum is taken over the top intersection numbers of all ample line bundles A occurring in a decomposition 2.3 on some modification of X . In fact, Fujita's proved the *upper* bound on $\text{Vol}(L)$ and the lower bound is considered to be substantially easier. The following two analytical versions are due to Boucksom [10]:

$$(4.20) \quad \text{Vol}(L) = \sup_{\phi' \in \mathcal{L}_{(X,L),a}} \int_{X - \{\phi' = -\infty\}} (dd^c \phi')^n / n! = \sup_{\phi' \in \mathcal{L}_{(X,L)}} \int_X ((dd^c \phi')_{ac})^n / n!$$

where $\mathcal{L}_{(X,L),a}$ denotes the subspace of metrics with analytical singularities and $((dd^c \phi')_{ac})$ denotes the absolutely continuous part of the current $dd^c \phi'$. The equivalence between 4.19 and the first equality in 4.20 is simply obtained by taking a log resolution of the pair $(X, \{\phi' = -\infty\})$ (compare [10]). The equivalence between 4.20 and corollary 4.8 is essentially contained in the proof of theorem 4.6: the lower bound on $\text{Vol}(L)$ follows from 4.13 and the upper bound from the fact that $(dd^c \phi_e)^n$ realizes the supremum in the last equality in 4.20 (and is approximated by ϕ_j in $\mathcal{L}_{(X,L),a}$).

Finally, a remark concerning the continuity of the volume function.

Remark 4.10. As is well-known the volume is continuous in the sense that for any line bundles L and F

$$\lim_{m \rightarrow \infty} m^{-n} \text{Vol}(mL + F) = \text{Vol}(L).$$

The continuity on the (open) cone of big line bundles is, for example, a simple consequence of the formula for Vol in corollary 4.8 (compare [10]). To get continuity up to the boundary of the big cone (for example that the limit is zero when L is non-big and pseudo-effective and F is ample) one can replace lemma 4.4 with the bound 4.13 (as in [10]). For a more direct algebro-geometric argument see [30] (I prop. 2.2.35).

4.1. The Bergman metric. The Hilbert space $\mathcal{H}(X, E(k))$ induces a metric on the line bundle L which may be expressed as

$$\phi_k(x) := k^{-1} \ln K_k(x, x) - k^{-1} \phi_F(x)$$

When $E(k) = L^k$ and $\phi_F = 0$ this metric is in the class $\mathcal{L}_{(X,L)}$ and is often referred to as the k th Bergman metric on L . If L is an ample line bundle, then this is the smooth metric on L obtained as the pull-back of the Fubini-Study metric on the hyperplane line bundle $\mathcal{O}(1)$ over $\mathbb{P}^N (= \mathbb{P}\mathcal{H}(X, L^k))$ (compare example 5.1 in section 5) under the Kodaira map

$$X \rightarrow \mathbb{P}\mathcal{H}(X, L^k), \quad y \mapsto (\Psi_1(x) : \Psi_2(x) : \dots : \Psi_N(x)) ,$$

for k sufficiently large, where (Ψ_i) is an orthonormal base for $\mathcal{H}(X, L^k)$ [23]. Note that the metric ϕ_k on L is singular precisely on the base locus $\text{Bs}(|E(k)|)$ and its curvature current is “almost positive” when k is large. The (almost positive) measures

$$1_{X-\text{Bs}(|E(k)|)}(dd^c(k^{-1}\ln K_k(x, x)))^n/n!$$

on X will be referred to as the k th Bergman volume forms.

Now we can prove the following theorem:

Theorem 4.11. *Let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, E(k))$. Then the following convergence holds:*

$$(4.21) \quad k^{-1}\phi_k \rightarrow \phi_e$$

uniformly on any fixed compact subset Ω of $X - \mathbb{B}_+(L)$. More precisely,

$$e^{-k(\phi - \phi_e)} C_\Omega^{-1} \leq B_k \leq C_\Omega k^n e^{-k(\phi - \phi_e)}$$

Moreover, the corresponding k th Bergman volume forms converge to the equilibrium measure:

$$(4.22) \quad 1_{X-\text{Bs}(|E(k)|)}(dd^c(k^{-1}\ln K_k(x, x)))^n/n! \rightarrow \mu_\phi$$

weakly as measures on X .

Proof. In the following proof it will be convenient to let C denote a sufficiently large constant (which may hence vary from line to line). First observe that when (F, ϕ_F) is trivial taking the logarithm of the inequality 4.6 in lemma 4.2 immediately gives the upper bound

$$k^{-1}\ln K_k(x, x) \leq \phi_e(x) + C \ln k/k$$

and the general case is completely analogous.

To get a lower bound, fix a point x_0 in $X - \mathbb{B}_+(L)$. By the extremal property 4.3 it is enough to find a section α_k in $\mathcal{H}(X, E(k))$ such that

$$(4.23) \quad |\alpha_k(x_0)|_{k\phi_e + \phi_F} \geq 1/C \quad \|\alpha_k\|_{X, k\phi_e + \phi_F} \leq C.$$

To this end take a “strictly positively curved modification” ψ_k of the metric $k\phi_e + \phi_F$ furnished by lemma 2.5. Then the extension theorem 2.4 gives a section α_k in $\mathcal{H}(X, E(k))$ such that

$$|\alpha_k(x_0)|_{\psi_k} \geq 1/C, \quad \|\alpha_k\|_{X, \psi_k} \leq C,$$

Applying 2.7 to the first inequality above and then 2.8 to the second one proves 4.23 (also using that by definition $\phi_e \leq \phi$).

To prove 4.22 first observe that the weak Monge-Ampere convergence 4.22 on the open set $X - \mathbb{B}_+(L)$ follows from the uniform convergence 4.21 (see [24]). Finally, by general integration theory it is (by theorem 3.4 (d)) enough to prove

$$\lim_{k \rightarrow \infty} \int_{X-\text{Bs}(|E(k)|)} (dd^c(k^{-1}\ln K_k(x, x)))^n/n! = \int_{X-\mathbb{B}_+(L)} (dd^c\phi_e)^n/n!$$

To this end first note that the weak Monge-Ampere convergence on $X - \mathbb{B}_+(L)$ implies

$$\liminf_k \int_{X - \text{Bs}(|E(k)|)} (dd^c(k^{-1} \ln K_k(x, x)))^n / n! \geq \int_{X - \mathbb{B}_+(L)} (dd^c \phi_e)^n n! /$$

Next, applying formula 4.20 to $\phi' = \ln K_k(x, x)$ shows that the left hand side (but with limsup instead) is bounded by $\text{Vol}(L)$ (compare remark 4.9). By corollary 4.18 this proves 4.22 and finishes the proof of the theorem. $\square$

For any line bundle L over X the intersection of the zero-sets of n “generic” sections in $H^0(X, L^k)$ with $X - \text{Bs}(|kL|)$ is a finite number of points (as follows from Bertini’s theorem [23]). The number of points is called the *moving intersection number* and is denoted by $(kL)^{[n]}$. The following corollary was obtained in [20] from Fujita’s theorem (see [30] for further references).

Corollary 4.12. *If L is a big line bundle then*

$$\text{Vol}(L) = \lim_{k \rightarrow \infty} \frac{(kL)^{[n]}}{k^n}$$

Proof. The proof follows immediately from (ii) in theorem 4.11 and the following fact:

$$n!(kL)^{[n]} = \int_{X - \mathbb{B}_k} (dd^c(\ln K_k(x, x)))^n.$$

The formula may be deduced by taking a log resolution of the pair $(X, \mathbb{B}_k(L))$ (compare [10]). But it also follows from properties of “random zeroes”, once one accepts that

$$n!(kL)^{[n]} = \int_{X - \mathbb{B}_k} Z_{f_1} \wedge \dots \wedge Z_{f_n},$$

where Z_{f_i} denotes the integration current determined by the zero-set of f_i , is independent of a “generic” tuple $(f_1, \dots, f_n)$ in $(H^0(X, L^k))^n$. Indeed, by proposition 2.2 in [36] the right hand side may be written as

$$\int_{X - \mathbb{B}_k} \mathbb{E}(Z_{f_1} \wedge \dots \wedge Z_{f_n}),$$

where $\mathbb{E}(Z_{f_1} \wedge \dots \wedge Z_{f_n})$ denotes the expectation value (taking values in the space of measures)⁸ of the intersection of the zero currents of n random independent sections in $\mathcal{H}(X, L^k)$. Changing the order of integration gives

$$\int_{X - \mathbb{B}_k} \mathbb{E}(Z_{f_1} \wedge \dots) = \mathbb{E} \int_{X - \mathbb{B}_k} (Z_{f_1} \wedge \dots) = n! \mathbb{E}(kL)^{[n]} = n!(kL)^{[n]}$$

$\square$

⁸ $\mathbb{E}(\cdot)$ denotes integration with respect to the Gaussian probability measure on the product $(\mathcal{H}(X, L^k))^n$ of Hilbert spaces.

4.2. The full Bergman kernel. Combining the convergence in theorem 4.6 with the local inequalities 4.5, gives the following convergence for the point-wise norm of the full Bergman kernel $K_k(x, y)$. The proof is completely analogous to the proof of theorem 2.4 in part 1 of [4].

Theorem 4.13. *Let L be a line bundle and let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, E(k))$. Then*

$$k^{-n} |K_k(x, y)|_{k\phi}^2 \omega_n(x) \wedge \omega_n(y) \rightarrow \Delta \wedge \mu_\phi ,$$

as measures on $X \times X$, in the weak $*$ -topology, where Δ is the current of integration along the diagonal in $X \times X$.

Finally, we will show that around any interior point of the set $D \cap X(0) - \mathbb{B}_+(L)$ the Bergman kernel $K_k(x, y)$ admits a complete local asymptotic expansion in powers of k , such that the coefficients of the corresponding symbol expansion coincide with the Tian-Zelditch-Catlin expansion (concerning the case when the curvature form of ϕ is positive on all of X ; see [8] and the references therein for the precise meaning of the asymptotic expansion). We will use the notation $\phi(x, y)$ for a fixed almost holomorphic-anti-holomorphic extension of a local representation of the metric ϕ from the diagonal Δ in $\mathbb{C}^n \times \mathbb{C}^n$, i.e. an extension such that the anti-holomorphic derivatives in x and the holomorphic derivatives in y vanish to infinite order along Δ .

Theorem 4.14. *Let L be a line bundle and let K_k be the Bergman kernel of the Hilbert space $\mathcal{H}(X, L^k)$. Any interior point in $D \cap X(0) - \mathbb{B}_+(L)$ has a neighbourhood where $K_k(x, y)e^{-k\phi(x)/2}e^{-k\phi(y)/2}$ admits an asymptotic expansion as*

$$(4.24) \quad k^n (\det(dd^c \phi)(x) + b_1(x, y)k^{-1} + b_2(x, y)k^{-2} + \dots) e^{k\phi(x, y)},$$

where b_i are global well-defined functions expressed as polynomials in the covariant derivatives of $dd^c \phi$ (and of the curvature of the metric ω) which can be obtained by the recursion given in [8].

Proof. The proof is obtained by adapting the construction in [8], concerning globally *positive* Hermitian line bundles, to the present situation. The approach in [8] is to first construct a “local asymptotic Bergman kernel” with the asymptotic expansion 4.24 close to any point where ϕ is smooth and $dd^c \phi > 0$. Hence, the local construction applies to the interior of the set $D \cap X(0) - \mathbb{B}_+(L)$ as well. Then the local kernel is shown to differ from the true kernel by a term of order $O(k^{-\infty})$, by solving a $\bar{\partial}$ -equation with a good L^2 -estimate. This is possible since $dd^c \phi > 1/C$ globally in that case. In the present situation we are done if we can solve

$$(4.25) \quad \bar{\partial} u_k = g_k,$$

where g_k is a $\bar{\partial}$ -closed $(0, 1)$ -form with values in L^k , supported on the interior of the bounded set $D \cap X(0) - \mathbb{B}_+(L)$ with an estimate

$$(4.26) \quad \|u_k\|_{k\phi+\phi_F} \leq C \|g_k\|_{k\phi+\phi_F}$$

To this end note that proceeding precisely as in step 2 in the proof of lemma 4.4, gives according to formula 4.12 a solution u_k satisfying 4.26, but with ϕ replaced with ϕ_e in the norm of g_k . However, since $\phi_e = \phi$ on the set where g_k is assumed to be supported this does prove 4.26 and hence finishes the proof of the theorem. $\square$

5. EXAMPLES

Finally, we illustrate some of the previous results with the following examples, which can be seen as variants of the setting considered in [5].

Example 5.1. Let X be the n -dimensional projective space $\mathbb{P}^n$ and let L be the hyperplane line bundle $\mathcal{O}(1)$. Then $H^0(X, L^k)$ is the space of homogeneous polynomials in of degree k in the $n + 1$ homogeneous coordinates $Z_0, Z_1, \dots, Z_n$. The Fubini-Study metric ϕ_{FS} on $\mathcal{O}(1)$ may be suggestively written as $\phi_{FS}(Z) = \ln(|Z|^2)$ and the Fubini-Study metric ω_{FS} on $\mathbb{P}^n$ is the normalized curvature form $dd^c \phi_{FS}$. Hence the induced norm on $H^0(X, L^k)$ is invariant under the standard action of $SU(n + 1)$ on $\mathbb{P}^n$. We may identify $\mathbb{C}^n$ with the “affine piece” $\mathbb{P}^n - H_\infty$ where H_∞ is the “hyperplane at infinity” in $\mathbb{C}^n$ (defined as the set where $Z_0 = 0$). In terms of the standard trivialization of $\mathcal{O}(1)$ over $\mathbb{C}^n$ (obtained by setting $Z_0 = 1$) the space $H^0(Y, L^k)$ may be identified with the space of polynomials $f_k(\zeta)$ in $\mathbb{C}_\zeta^n$ of total degree at most k and the metric ϕ_{FS} on $\mathcal{O}(1)$ may be represented by the function

$$\phi_{FS}(\zeta) = \ln(1 + |\zeta|^2).$$

Moreover, any smooth metric on $\mathcal{O}(1)$ may be represented by a function $\phi(\zeta)$ satisfying the following necessary growth condition⁹

$$(5.1) \quad -C + \ln(1 + |\zeta|^2) \leq \phi(\zeta) \leq \ln(1 + |\zeta|^2) + C,$$

which makes sure that the norm 1.9, expressed as

$$\|f_k\|_{k\phi}^2 := \int_{\mathbb{C}^n} |f_k(\zeta)|^2 e^{-k\phi(\zeta)} \omega_{FS}^n / n!$$

is finite when f_k corresponds to a section of the k th power of $\mathcal{O}(m)$, for $m = 1$. In particular, any smooth compactly supported function $\chi(\zeta)$ determines a *smooth* perturbation

$$(5.2) \quad \phi_\chi(\zeta) := \phi_{FS}(\zeta) + \chi(\zeta)$$

of ϕ_{FS} on $\mathcal{O}(1)$ over $\mathbb{P}^n$, to which the results in section 3 and 4 apply.

The next class of example is offered by toric varieties.

Example 5.2. Let Δ be a Delzant polytope in $\mathbb{R}^n$ obtained as the convex hull of points in $\mathbb{Z}^n$ (see [1]) It induces a triple $(X_\Delta, L_\Delta, \phi_\Delta)$, where X_Δ , is an n -dimensional complex compact projective manifold on which the complex torus $\mathbb{C}^{*n}$ acts effectively with an open dense orbit and (L_Δ, ϕ_Δ)

⁹in order that ϕ extend over the hyperplane at infinity to a *smooth* metric further conditions are needed.

is an Hermitian positive line bundle, invariant under the action of T^n (the real torus in $\mathbb{C}^{*n}$). The curvature form $dd^c\phi$ defines a T^n -invariant Kähler metric on X_Δ . Identifying $\mathbb{C}_z^{*n}$ with an open dense set in X_Δ , the space $H^0(X_\Delta, L_\Delta^k)$ may be identified with the space spanned by all monomials z^α with α a multi-index in the scaled polytope $k\Delta$ and the metric ϕ_Δ may be identified with a plurisubharmonic function on $\mathbb{C}_z^{*n}$:

$$\phi_\Delta(z) = \ln\left(\sum_{\alpha \in \Delta \cap \mathbb{Z}^n} |z^\alpha|^2\right).$$

Writing $v_i := \ln(|z_i|^2)$ identifies $\phi_\Delta(z)$ with a convex function on $\mathbb{R}^n$ that we, by a slight abusive of notation, denote by $\phi_\Delta(v)$. Now any real smooth compactly supported function $\chi(v)$ on $\mathbb{R}^n$ induces a perturbation $\phi := \phi_\Delta + \chi$, yielding a new smooth T^n -invariant metric ϕ on L_Δ . In this notation the almost everywhere convergence of the Bergman function B_k (theorem 4.6) may be written as

$$\sum_{p \in \Delta \cap (\frac{1}{k}\mathbb{Z})^n} \frac{e^{k(\langle p, v \rangle - \phi(v))}}{\int_{v \in \mathbb{R}^n} e^{k(\langle p, v \rangle - \phi(v))} \det(\frac{\partial^2 \phi_\Delta}{\partial^2 v}) dv} \rightarrow 1_D(v) \left(\frac{2}{\pi}\right)^n \frac{\det(\frac{\partial^2 \phi}{\partial^2 v})(v)}{\det(\frac{\partial^2 \phi_\Delta}{\partial^2 v})(v)}$$

where $\langle p, v \rangle$ and dv denote the Euclidian scalar product and volume form, respectively.

Note that in the logarithmic coordinates v_i we have that $(\frac{\partial^2 \phi}{\partial z_i \partial \bar{z}_j}) = (\frac{\partial^2 \phi}{\partial v_i \partial v_j})/4$. Hence, it follows (more or less from the definition) that the graph of the equilibrium metric ϕ_e determined by ϕ is simply the *convex hull* of the graph of ϕ considered as a function of v . In particular, in the “generic” toric case ϕ_e will not be in the class $\mathcal{C}^2$. Indeed, consider for example the case when $n = 1$, so that $X_\Delta = \mathbb{P}^1$, and take $\phi(v)$ to be an even function with two non-degenerate minima at $\pm a$ (figure 1.1 in section 1). Then $\frac{\partial^2 \phi}{\partial^2 v}(a) > 0$, but $\frac{\partial^2 \phi_e}{\partial^2 v}(a - \epsilon) = 0$ if $0 < \epsilon < 2a$. Note that if we deform ϕ in such a way that ϕ_e is unchanged (figure 5.1), then the leading Bergman kernel asymptotics for the Bergman function B_k given by theorem 4.6 are also unchanged. However, the exponential decay of B_k in the complement of the set D does change. For example, by theorem 4.11:

$$\lim_{k \rightarrow \infty} \inf_X (B_k)^{1/k} = e^{-\sup_X (\phi_e - \phi)},$$

i.e. minus the exponential of the limit equals the supremum of the distance between points on the graph of ϕ and points on the graph of ϕ_e . In [24] the limit in the left hand side above is called the *Tchebishev constant* and the right hand side the *Alexander capacity* (associated to the tripple (X, L, ϕ)).

The following basic example of a big (non-ample) line bundle shows that ϕ_e may be singular on all of $\mathbb{B}_+(L)$:

Example 5.3. Let X be the blow-up of $\mathbb{P}^2$ and denote by π the projection (blow-down map) from X to $\mathbb{P}^2$. Let $L = \pi^*\mathcal{O}(1) \otimes [E]$, where E is the




FIGURE 5.1. The picture represents the graph of a metric ϕ on $\mathcal{O}(2)$ over $\mathbb{P}^1$, in logarithmic radial coordinates, which is a compact perturbation ($\phi = \phi_\Delta + \chi$) of the Fubini-Study metric $\phi_\Delta(v) (= \ln(e^{-v} + e^v) + C)$. Deforming ϕ without changing the “convex hull” ϕ_e does not change the leading Bergman function asymptotics.

exceptional divisor. Since, $\int_E c_1(L) < 0$ any element of $\mathcal{L}_{(X,L)}$ is identically equal to $-\infty$ on E . Moreover, as is well-known $E = \mathbb{B}_+(L)$. In particular, $\phi_e \equiv -\infty$ on $\mathbb{B}_+(L)$.

REFERENCES

- [1] Abreu, M: Kahler geometry of toric manifolds in symplectic coordinates. Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montreal, QC, 2001), 1–24, Fields Inst. Commun., 35, Amer. Math. Soc., Providence, RI, 2003.
- [2] Bedford, E; Taylor, A: The Dirichlet problem for a complex Monge-Ampere equation. Invent. Math 37 (1976), no 1, 1-44
- [3] Berman, R: Bergman kernels and local holomorphic Morse inequalities. Math Z., Vol 248, Nr 2 (2004), 325–344
- [4] Berman, R: Super Toeplitz operators on holomorphic line bundles J. Geom. Anal. 16 (2006), no. 1, 1–22.
- [5] Berman, R: Bergman kernels and weighted equilibrium measures of $\mathbb{C}^n$. Preprint in 2007 at arXiv.org/abs/math.CV/0702357
- [6] Berman, R: Bergman kernels and equilibrium measures for ample line bundles. Preprint in 2007 at [arXiv:0704.1640](http://arXiv.org/abs/math.CV/0704.1640)
- [7] Berman, R: Bergman kernels and equilibrium measures vanishing along global multiplier ideal sheaves. (in preparation)
- [8] Berman R; Berndtsson B; Sjöstrand J: Asymptotics of Bergman kernels. Preprint in 2005 at arXiv.org/abs/math.CV/050636.
- [9] Berndtsson, Bo: Positivity of direct image bundles and convexity on the space of Kähler metrics. Preprint in 2006 at arXiv.org/abs/math.CV/0608385
- [10] Boucksom, S: On the volume of a line bundle. Internat. J. Math 13 (2002), no 10. 1043-1063
- [11] Boucksom, S: Divisorial Zariski decompositions on compact complex manifolds. Ann. Sci. École Norm. Sup. (4) 37 (2004), no. 1, 45–76.

- [12] Bremermann, H: On the conjecture of equivalence between the plurisubharmonic functions and the Harthogs functions. *Math. Ann.* 131 (1956), 76-86
- [13] Chen, X., The space of Kähler metrics, *J. Differential Geom.* 56 (2000), no. 2, 189-234.
- [14] Chen, X; Tian, G: Geometry of Kähler metrics and foliations by discs. Preprint in 2004 at arXiv.org/abs/math.DG/0409433.
- [15] Demailly, J-P: Champs magnetiques et inegalite de Morse pour la d"-cohomologie., *Ann Inst Fourier*, 355 (1985), 185-229
- [16] Demailly, J-P: Complex analytic and algebraic geometry. Available at www-fourier.ujf-grenoble.fr/~demailly/books.html
- [17] Demailly, J-P: Potential Theory in Several Complex Variables. Manuscript available at www-fourier.ujf-grenoble.fr/~demailly/
- [18] Demailly, J-P: Regularization of closed positive currents and Intersection theory. *J. Alg. Geom.* 1 (1992), 361-409
- [19] Demailly, J-P: Multiplier ideal sheaves and analytic methods in algebraic geometry. School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000), 1-148, ICTP Lect. Notes, 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.
- [20] Demailly, J-P; Ein, L; Lazarsfeld, R A subadditivity property of multiplier ideals. *Michigan Math. J.* 48 (2000)
- [21] Ein, L; Lazarsfeld, R; Mustata, M; Nakamaye, M; Popa, M Restricted volumes and base loci of linear series. Preprint in 2006 at arXiv:math/0607221
- [22] Fujita, T: Approximating Zariski decomposition of big line bundles. *Kodai Math. J.* 17 (1994), no. 1, 1-3.
- [23] Griffiths, P; Harris, J: Principles of algebraic geometry. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1994.
- [24] Guedj, V; Zeriahi, A: Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.* 15 (2005), no. 4, 607-639.
- [25] Hedenmalm, H; Makarov, N: Quantum Hele-Shaw flow, Preprint in 2004 at arXiv.org/abs/math.PR/0411437
- [26] Keller, J; Szekelyhidi, G; Thomas, R: "Bergman kernels and slope stability" (work in progress).
- [27] Kinderlehrer, D; Stampacchia: An introduction to variational inequalities and their applications, Academic Press, New York, 1980
- [28] Klimek, M: Pluripotential theory. London Mathematical Society Monographs. New Series, 6. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1991
- [29] Kolodziej, S: The complex Monge-Ampere equation and pluripotential theory. *Mem. Amer. Math. Soc.* 178 (2005), no. 840
- [30] Lazarsfeld, R: Positivity in algebraic geometry. I. Classical setting: line bundles and linear series. II. Positivity for vector bundles, and multiplier ideals. A Series of Modern Surveys in Mathematics, 48. and 49. Springer-Verlag, Berlin, 2004.
- [31] Phong, D.H; Sturm, J: The Monge-Ampere operator and geodesics in the space of Kähler potentials. *Invent. math.* 166 (2006), 125-149
- [32] Saff, E; Totik, V: Logarithmic potentials with exterior fields. Springer-Verlag, Berlin. (1997) (with an appendix by Bloom, T)
- [33] Shiffman, B; Zelditch, S: Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Comm. Math. Phys.* 200 (1999), no. 3, 661-683.
- [34] Shiffman, B; Zelditch, S: Equilibrium distribution of zeros of random polynomials. *Int. Math. Res. Not.* 2003, no. 1, 25-49.
- [35] Shiffman, B; Zelditch, S: Random polynomials with prescribed Newton polytope. *J. Amer. Math. Soc.* 17 (2004), no. 1, 49-108 (electronic).

- [36] Shiffman, B; Zelditch S: Number variance of random zeros on complex manifolds. Prinprint in 2006 at arXiv:math/0608743 (to appear in GAFA)
- [37] Zelditch, S: Szegő kernels and a theorem of Tian. Internat. Math. Res. Notices 1998, no. 6, 317–331.

E-mail address: robertb@math.chalmers.se

Current address: Institut Fourier, 100 rue des Maths, BP 74, 38402 St Martin d'Heres (France)